

Aberystwyth University

Hydrology of debris-covered glaciers in High Mountain Asia

Miles, Katie E.; Hubbard, Bryn; Irvine-fynn, Tristram D.I.; Miles, Evan S.; Quincey, Duncan J.; Rowan, Ann V.

Published in:
Earth-Science Reviews

DOI:
[10.1016/j.earscirev.2020.103212](https://doi.org/10.1016/j.earscirev.2020.103212)

Publication date:
2020

Citation for published version (APA):

Miles, K. E., Hubbard, B., Irvine-fynn, T. D. L., Miles, E. S., Quincey, D. J., & Rowan, A. V. (2020). Hydrology of debris-covered glaciers in High Mountain Asia. *Earth-Science Reviews*, 207, [103212].
<https://doi.org/10.1016/j.earscirev.2020.103212>

Document License CC BY-NC-ND

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Hydrology of debris-covered glaciers in High Mountain Asia

Author's post-print/accepted version
(CC-BY-NC-ND)

DOI of published version (Earth-Science Reviews):

<https://doi.org/10.1016/j.earscirev.2020.103212>

Hydrology of debris-covered glaciers in High Mountain Asia

Katie E. Miles^{1*}, Bryn Hubbard¹, Tristram D. L. Irvine-Fynn¹, Evan S. Miles², Duncan J. Quincey³
and Ann V. Rowan⁴

1. Centre for Glaciology, Department of Geography and Earth Sciences, Aberystwyth University, Aberystwyth, UK

2. Swiss Federal Research Institute WSL, 8903 Birmensdorf, Switzerland

3. School of Geography, University of Leeds, Leeds, UK

4. Department of Geography, University of Sheffield, Sheffield, UK

** Correspondence to: kam64@aber.ac.uk*

Key words

Glaciers; debris-covered glaciers; glacier hydrology; High Mountain Asia

Abstract

The hydrological characteristics of debris-covered glaciers are known to be fundamentally different from those of clean-ice glaciers, even within the same climatological, geological, and geomorphological setting. Understanding how these characteristics influence the timing and magnitude of meltwater discharge is particularly important for regions where downstream communities rely on this resource for sanitation, irrigation, and hydropower, as in High Mountain Asia. The hydrology of debris-covered glaciers is complex: rugged surface topographies typically route meltwater through compound supraglacial-englacial systems involving both channels and ponds, as well as pathways that remain unknown. Low-gradient tongues that extend several kilometres retard water conveyance and promote englacial storage. Englacial conduits are frequently abandoned and reactivated as water supply changes, new lines of permeability are exploited, and drainage is captured due to high rates of surface and subsurface change. Seasonal influences, such as the monsoon, are superimposed on these distinctive characteristics, reorganising surface and subsurface drainage rapidly from one season to the next. Recent advances in understanding have mostly come from studies aimed at quantifying and describing supraglacial processes; little is known about the subsurface hydrology, particularly the nature (or even existence) of subglacial drainage. In this review, we consider in turn the supraglacial, englacial, subglacial, and proglacial hydrological domains of debris-covered glaciers in High Mountain Asia. We summarise different lines of evidence to establish the current state of knowledge and, in doing so, identify major knowledge gaps. Finally, we use this information to suggest six themes for future hydrological research at High Mountain Asian debris-covered glaciers in order to make timely long-term predictions of changes in the water they supply.

36 1. Introduction

37 Debris-covered glaciers have gained increased research attention over recent years, partly in
 38 recognition of their role as water sources for large parts of the world's population (Immerzeel et
 39 al., 2020; Scherler et al., 2011) and partly because they host a range of distinctive features, driven
 40 by processes that are largely absent from their clean-ice counterparts. Definitions of what
 41 constitutes a 'debris-covered glacier' vary widely (e.g. Anderson, 2000; Kirkbride, 2011), but here
 42 we define them to be glaciers with a largely continuous layer of supraglacial debris over most of
 43 the ablation area, typically increasing in thickness towards the terminus (Figure 1). Debris can be
 44 supplied to such glaciers by snow avalanches, rockfalls, and landslides from local mountainsides
 45 onto the glacier surface (Figure 2, 3A), melt-out of englacial debris, thrusting transporting debris
 46 from the glacier bed, dust blown from exposed moraines, or solifluction from (ice-cored) moraines
 47 (Dunning et al., 2015; Gibson et al., 2017b; Hambrey et al., 2008; Kirkbride and Deline, 2013;
 48 Rowan et al., 2015; van Woerkom et al., 2019). The surface debris layer can range in thickness
 49 from scattered particles to several metres, including large rocks and substantial boulders (Figure
 50 3C and D) (Inoue and Yoshida, 1980; McCarthy et al., 2017; Nicholson et al., 2018).




Figure 1 – Debris-covered glaciers in the Sagarmatha National Park, Nepal Himalaya, annotated with some of the features distinctive to High Mountain Asian debris-covered glaciers. **A)** Chola Glacier (image width is ~1.5 km across the glacier terminus and lake). **B)** Imja-Lhotse Shar Glacier, showing the terminus and calving front (~0.75 km width) into Imja Tsho, looking towards the accumulation area of the tributary Amphulapcha Glacier. **C)** Khumbu Glacier, showing the upper ablation area (clean-ice flowing from the Khumbu Icefall) to the left and the ~8 km long lower

ablation area (debris-covered tongue) to the right; dashed yellow lines are 100 m contours. Image credit for A and B: Katie Miles; and C: Tristram Irvine-Fynn.

Debris-covered glaciers are present in nearly all of Earth's glacierised regions, with a particularly large concentration in High Mountain Asia (Bolch et al., 2012; Kraaijenbrink et al., 2017; Scherler et al., 2018); sub-regional variability in the debris cover of which is presented in Brun et al. (2019) (their Figure 1). Debris-covered glaciers therefore contribute an important proportion of streamflow used for drinking water, irrigation, and hydroelectric power; this streamflow is particularly effective in reducing seasonal water shortages (Bolch et al., 2019; Immerzeel et al., 2020, 2010; Pritchard, 2019; Scott et al., 2019). Glacier mass loss in response to climate warming is currently increasing river discharge and contributions to sea level (Hock et al., 2019; Lutz et al., 2014; Radić et al., 2014; Shea and Immerzeel, 2016), but studies simulating future scenarios universally project long-term reductions in flow, perhaps as soon as 2050 in central Asia (Barnett et al., 2005; Bolch et al., 2012; Huss and Hock, 2018; Lutz et al., 2014; Ragettli et al., 2016b; Rounce et al., 2020; Sorg et al., 2012). Passing of 'peak water' threatens future water security in many regions, particularly across High Mountain Asia (Bolch et al., 2019; Eriksson et al., 2009; Hannah et al., 2005; Huss and Hock, 2018; Immerzeel et al., 2010; Winiger et al., 2005). A decrease in discharge from the Indus and Brahmaputra rivers alone is estimated to affect 260 million people (Immerzeel et al., 2010).

The long-term response of debris-covered glaciers to changing climatic conditions is non-linear and results from complexities relating to spatial variability in debris concentration and climatic controls integrated over at least several decades (Benn et al., 2012; Vaughan et al., 2013). A multidecadal trend of surface lowering, stagnation, and glacier mass loss has already been observed on many debris-covered glaciers across High Mountain Asia (Bolch et al., 2012, 2011; Brun et al., 2017; Dehecq et al., 2019; Hock et al., 2019; Kääb et al., 2012; Pellicciotti et al., 2015; Scherler et al., 2011) as a result of warmer air temperatures and weaker monsoons (Pieczonek et al., 2013; Thakuri et al., 2014). However, predictions of mass loss from individual glacierised regions vary considerably. For example, in the Everest region of the Himalaya, estimates of ice mass loss by 2100 vary from ~10% (Rowan et al., 2015), through 50% (Soncini et al., 2016), to 99% in extreme scenarios (warming of ~3°C) (Shea et al., 2015). Model outputs also vary spatially at a regional scale (e.g. Chaturvedi et al., 2014; Kraaijenbrink et al., 2017; Zhao et al., 2014). Such projections depend on the future climate scenario used, but a number of key knowledge gaps also exist concerning the character of debris-covered glaciers and the processes influencing their varied geometrical response to climate change (Benn et al., 2012; Bolch et al., 2012; Huss, 2011; Scherler et al., 2011).

Understanding how meltwater is produced, transported, and stored within High Mountain Asian debris-covered glaciers is therefore imperative. There is growing recognition that the configuration and efficiency (i.e. bulk system transit velocity) of water routing across and through debris-covered ice is distinctively different from that of clean-ice glaciers, even within the same glacial system. This was first shown by a recent study on Miage Glacier, a debris-covered glacier in the Italian Alps (Fyffe et al., 2019b). Debris-covered glacier surfaces are complex, particularly those in High Mountain Asia, the ablation areas of which are often characterised by hummocky, rugged

topography atop a shallow (or even reversed) longitudinal surface gradient (Figures 1 and 2). This commonly results from an inverted mass-balance regime, where the greatest ablation rates are experienced in the middle, rather than lower, ablation area (King et al., 2017). Debris-covered ablation areas also exhibit bare ice cliffs and supraglacial ponds – depressions capable of storing meltwater for both short and long periods within nested catchments of varying spatial scales (Section 2) – and these glaciers frequently terminate in proglacial lakes (Section 5). Other unique characteristics of High Mountain Asian debris-covered glaciers include the accumulation areas often being at extremely high elevations, with a steep surface gradient (often an icefall) transporting ice into the ablation area (Figure 1). These features provide a setting that strongly influences the nature of hydrological systems in this region (Benn et al., 2017; Miles et al., 2019), but has restricted hydrological research due to the remoteness and inaccessibility of such glaciers.

In this review, we consider the current state of knowledge of debris-covered glacier hydrological systems in High Mountain Asia. Four hydrological domains are considered in turn: supraglacial (Section 2), englacial (Section 3), subglacial (Section 4), and proglacial (Section 5). Within each section, we summarise existing research and understanding of debris-covered glacier hydrological systems and then address key remaining knowledge gaps. Figure 2 provides a reference conceptual diagram of a (High Mountain Asian) debris-covered glacier, with each hydrological feature encompassing both known and unknown elements of each domain. Finally, in light of the review, we propose six future research themes concerning the hydrology of debris-covered glaciers (Section 6). This review is intended to complement existing reviews of clean-ice valley glacier hydrology (e.g. Fountain and Walder, 1998; Hubbard and Nienow, 1997; Irvine-Fynn et al., 2011; Jansson et al., 2003). We note that there are a number of differing climatic regimes across High Mountain Asia, with precipitation in particular varying closely with topography (Bookhagen and Burbank, 2006); these climatic regimes will influence the thermal regime, geometry, mass balance, and thus hydrology of the glaciers in each of these sub-regions. While our review draws on research carried out across High Mountain Asia, much of that research has been carried out in the monsoon-influenced Himalaya, particularly Nepal, from where the review and our illustrations of many of the key elements draw strongly.




Figure 2 – A conceptual illustration of the main landscape and hydrological features of a typical debris-covered glacier. Features specific to debris-covered glaciers in High Mountain Asia are labelled in Figure 1.


2. Supraglacial hydrology

2.1 Supraglacial zone

2.1.1 Meltwater generation

Supraglacial meltwater is produced on debris-covered glaciers through ablation of surface ice and snow, with the spatial pattern of melt complicated by the surface debris extent, thickness, and lithological characteristics (Figures 1 and 3). A debris layer shallower than the critical thickness, typically ~ 0.05 m, decreases albedo and thus increases the ablation rate compared to debris-free ice (Figure 4). The ablation rate peaks at a debris thickness of ~ 0.02 – 0.05 m, known as the effective thickness (Adhikary et al., 2000; Evatt et al., 2015; Inoue and Yoshida, 1980; Juen et al., 2014; Kayastha et al., 2000; Lejeune et al., 2013; Nicholson and Benn, 2013, 2006; Østrem, 1959; Singh et al., 2000; Takeuchi et al., 2000). The exact values of the critical and effective thickness strongly depend on the debris thermal conductivity, which can vary widely both across a glacier surface and in time according to whether the debris is wet or dry (Casey et al., 2012; Collier et al., 2015, 2014; Gibson et al., 2017b; Nicholson and Benn, 2013; Pelto, 2000). In contrast, a debris layer thicker than the critical thickness of ~ 0.05 m insulates the ice from incoming solar radiation, inhibiting the receipt of surface energy at the ice-debris interface and thus reducing the melt rate (Figure 4). Beneath a debris thickness of 0.25 – 0.30 m, ice becomes almost fully insulated from

daily surface energy fluxes, with only longer-term changes in surface energy balance reaching the underlying debris-ice interface (Bocchiola et al., 2015; Brock et al., 2010; Conway and Rasmussen, 2000; Nicholson and Benn, 2013; Østrem, 1959; Reid and Brock, 2010). In addition, turbulent energy fluxes have been shown to reduce net radiative fluxes at the debris surface (of a 0.75 m thick debris layer) by 17% over a full melt season, further diminishing the energy available for melt at the ice-debris interface (Steiner et al., 2018b). Variations in ablation according to these factors represent an important first-order control on glacier surface morphology and are partially responsible for the characteristic hummocky topography superimposed on a shallow or concave-upward (reversed gradient) debris-covered glacier surface profile (Figure 1).



*Figure 3 – Images illustrating debris transport processes, englacial debris inclusions, and variations in supraglacial debris thickness on Khumbu Glacier, Nepal Himalaya: **A**) a landslide scar (yellow circle, ~500 m wide) and unstable rock faces (purple circle) providing debris to the glacier surface; image is taken looking east across the surface of Khumbu Glacier, and the debris layer above ice cliffs can also be seen. **B**) an ice cliff with entrained debris (green circle), debris-rich ice layers (orange circle), and a moderately thick (~1–2 m) surface debris layer; **C**) a thin (~0.20 m; red arrow) surface debris layer above ice adjacent to a supraglacial pond; and **D**) a thick (> 5 m; red arrow) surface debris layer above an ice cliff. Image credit for A: Duncan Quincey; and B–D: Katie Miles.*

Counteracting the influence of a thick surface debris layer, the ablation rate of debris-covered glaciers is enhanced by the presence of supraglacial ponds (Section 2.1.2) and ice cliffs

(Figure 3B and D). The latter form by slumping of debris from steep slopes, calving at supraglacial pond margins (Section 2.1.2), or the collapse of englacial voids (Section 3.1), all of which expose steep, bare ice (Figure 3B) or thinly debris-covered faces (Figure 3D) at the glacier surface (Benn et al., 2012, 2001; Sakai et al., 2002; Thompson et al., 2016). The melting of ice cliffs can be responsible for a substantial proportion of debris-covered glacier ablation (Brun et al., 2016; Buri et al., 2016b; Han et al., 2010; Juen et al., 2014; Reid and Brock, 2014; Sakai et al., 2002, 2000; Thompson et al., 2016), accounting for 23–69% of the total ablation of debris-covered areas whilst covering a small proportion of the total glacier area. The ice cliffs exhibit melt rates that are 3–14 times higher than beneath debris-covered ice (Brun et al., 2018; Immerzeel et al., 2014; Sakai et al., 1998). Where ice cliffs are associated with supraglacial ponds, there is further potential for increased melting through undercutting and calving processes (Brun et al., 2016; Buri et al., 2016a; Miles et al., 2016; Röhl, 2008; Thompson et al., 2016). Taken together, ice cliff and pond systems may contribute considerably to the surface lowering of debris-covered glaciers in the central ablation area (King et al., 2017; Nuimura et al., 2012; Pellicciotti et al., 2015; Ragettli et al., 2016a; Thompson et al., 2016; Watson et al., 2017), contributing to the inverted mass-balance regime typical of High Mountain Asian debris-covered glaciers.




Figure 4 – Østrem curve examples showing variations in the relationship between debris thickness and ice ablation on different glaciers. (a) notes the effective thickness, namely the debris thickness at which maximum melt occurs. (b) marks the critical thickness, the debris thickness at which melt becomes inhibited compared to that of clean ice on different glaciers (indicated on both for Isfjallsglaciaren). From Nicholson & Benn (2006).

2.1.2 Meltwater storage

Supraglacial ponds (Figure 5), a term used here to include larger water bodies elsewhere sometimes referred to as lakes, are common and important features on debris-covered glaciers. Ponds are generally absent from clean-ice valley glaciers but are prevalent on low-gradient areas of ice sheet margins (Chu, 2014; Sundal et al., 2009). Similarly for debris-covered glaciers, the most important control on the location of supraglacial pond formation is a low glacier surface slope (Miles et al., 2017b; Quincey et al., 2007; Reynolds, 2000; Sakai, 2012; Sakai et al., 2000; Sakai and

Fujita, 2010; Salerno et al., 2012). A surface gradient of $\leq 2^\circ$ is considered to promote the development of larger ponds, while smaller isolated and transient ponds are considered more likely on steeper slopes (Miles et al., 2017b; Quincey et al., 2007; Reynolds, 2000). The upglacier slope has also been shown to have an influence, being inversely correlated to the total area of lakes downglacier (Salerno et al., 2012).

Glacier velocity and motion type also exert controls over supraglacial pond location. An increase in lake concentration is common towards the termini of debris-covered glaciers, areas that are typically characterised by low surface velocities (Kraaijenbrink et al., 2016b; Miles et al., 2017b; Quincey et al., 2007; Sakai, 2012; Salerno et al., 2015, 2012). A decrease in velocity towards both the glacier terminus and ice inflow at the confluences of flow units (Kraaijenbrink et al., 2016b) causes compressive flow, which tends to close crevasses and drive water back to the surface, as well as limiting effective drainage from the glacier surface (Kraaijenbrink et al., 2016b; Miles et al., 2017b). The thinning and stagnation of debris-covered glacier termini may also enhance meltwater production, further promoting the formation of ponds (Salerno et al., 2015; Thakuri et al., 2016).




Figure 5 – Examples of supraglacial pond size and temporal changes on Khumbu Glacier, Nepal Himalaya. Ponds range in diameter from: **A)** several metres; **B)** tens of metres (person circled in red for scale); **C)** and **D)** hundreds of metres; **E)** and **F)** several kilometres. A) and B) are located in the upper ablation area. C) and D) show the same pond-cliff-cave system in the mid-ablation area two years apart, with notable expansion of the cave via undercutting and calving. The pond, which has reduced in area (likely partly drained), was filled with a large amount of small, calved ice blocks in May 2019 and large cracks in the cliff system suggest further imminent large-scale calving. E) and F) show the expanding linked supraglacial pond chain at the terminus, also two years apart (green star indicates the same location as images were taken from slightly different positions). Pond growth and coalescence has progressively eroded the hummocks that formerly separated these

ponds. Higher melt rates are indicated by the covering of ice cliffs in fine debris ('dirty ice'). Image credit for A–D and F: Katie Miles; and E: Evan Miles.

Initial supraglacial pond growth occurs primarily through subaqueous melting at the base of any slight depression (Chikita et al., 1998; Mertes et al., 2016; Miles et al., 2016; Stokes et al., 2007; Thompson et al., 2012). Water accumulates and is heated by incoming solar radiation, causing the pond to warm. For example, Chikita et al. (1998) measured a maximum temperature of $\sim 5^{\circ}\text{C}$ at a supraglacial pond surface on Trakarding Glacier, Nepal Himalaya. Excess energy is thus available for lateral and vertical ablation wherever pond water is in contact with ice, increasing the pond size, steepening marginal slopes and mobilising debris to expose bare ice (Figure 5E and F) (Stokes et al., 2007). Subaqueous pond melt rates are greatest when bare ice is exposed or covered in a thin layer of debris; layers of thick sediment at the base of ponds effectively terminate bottom deepening by preventing transfer of energy from the warmer pond water to the ice surface (Horodyskyj, 2015). Furthermore, mixing of pond stratification by inflowing meltwater on Koxkar Glacier, Tien Shan, has been shown to increase the temperature (by $\sim 4^{\circ}\text{C}$) and density of the pond (Wang et al., 2012). Here, the warmed surface water sinks to the pond base and increases the potential for subaqueous melting; a process that can also be induced by wind-driven currents (Chikita et al., 1998).

Supraglacial ponds surrounded by ice cliffs tend to be larger and deeper than those without cliffs (Watson et al., 2018), as the ice cliffs facilitate pond growth by subaerial melting and backwasting, particularly during the monsoon melt season (Röhl, 2008; Steiner et al., 2019). Where warm surface pond water meets glacier ice, it can undercut the cliff beneath the waterline; progressive undercutting and thermo-erosional notch development may then lead to calving of the ice cliff and pond expansion (Figure 5C and D) (Chikita et al., 1998; Kirkbride and Warren, 1997; Mihalcea et al., 2006; Miles et al., 2016; Röhl, 2008, 2006; Sakai et al., 2009). Conversely, where the subaqueous and ice cliff melt rates are similar, the ice cliff will persist and backwaste stably (Brun et al., 2016; Buri et al., 2016a; Miles et al., 2016). Calving is most effective at larger ponds (Röhl, 2008), in particular where the fetch is greater than 20 m and the water temperature is $2\text{--}4^{\circ}\text{C}$ (Sakai et al., 2009). Calving events cause further mixing of pond layers, driving warmer surface water towards the base and again enhancing basal melting; the greatest supraglacial pond deepening rates have been shown to occur adjacent to the tallest calving ice cliffs (Thompson et al., 2012). Although sedimentation from ice cliffs and inflowing water can reduce pond depth, this effect is commonly outstripped by ablation (Thompson et al., 2012), resulting in general long-term pond growth.

A pattern of supraglacial pond evolution into ice-marginal moraine-dammed lakes has been observed for some ponds on debris-covered glaciers in High Mountain Asia. Supraglacial ponds form initially as 'perched ponds', isolated above the englacial drainage network (Benn et al., 2012). As these ponds increase in area and depth, they evolve from perched to base-level features, where the base-level is determined by the height at which water leaves the glacial system (usually the elevation of a spillway through the terminal moraine or the glacier bed, if water is transported there) (Mertes et al., 2016; Thompson et al., 2012; Watanabe et al., 2009). However, differing sub-catchments may have differing base-levels defined by other hydrological features such as moulins,

resulting in a stepped hydrological cascade based on several local base-levels. Alternatively, the presence of a groundwater system can result in a regional base-level. Over an extended period of glacier recession, an increasing number of supraglacial ponds form and grow over time, creating a chain of terminus-base-level ponds that eventually coalesce (Figure 5E and F) (Sakai, 2012; Salerno et al., 2012). The growth of base-level ponds is not limited by periodic drainage, potentially allowing dramatic increases in area, particularly through calving (Benn et al., 2001; Sakai, 2012; Thompson et al., 2012). If meltwater cannot escape from the system, pond expansion and coalescence may eventually lead to the formation of a single base-level moraine-dammed proglacial lake at the glacier terminus (Section 5.1.1) (Mertes et al., 2016; Watanabe et al., 2009) that will continue to expand both upglacier and downwards by ice melt.

Various stages of this supraglacial pond evolution are simultaneously present on many High Mountain Asian debris-covered glaciers. An increase in supraglacial pond area and proglacial lake formation, assumed to be in response to a warmer climate and glacier surface lowering, has been observed in recent decades in, for example, the Tien Shan (Wang et al., 2013), Bhutan Himalaya (Ageta et al., 2000; Komori, 2008), and Nepal Himalaya (Benn et al., 2000; Watson et al., 2016). Within the Hindu-Kush Himalaya, a clear divide has appeared between the East, where there are a greater number of larger ponds that have grown between 1990–2009 and become increasingly proglacial, and the West, where already generally smaller supraglacial ponds have been decreasing further in area (Gardelle et al., 2011). However, local variations do occur and the pattern is not universal (e.g. Steiner et al., 2019).

As isolated perched ponds grow, they can deepen such that they become connected to the englacial system by intersecting englacial flow pathways, and drain (Benn et al., 2001; Liu et al., 2015; Röhl, 2008; Watson et al., 2018, 2016; Wessels et al., 2002), temporarily halting further pond expansion (Mertes et al., 2016). Pond drainage is promoted in zones of higher local surface velocity and strain rates, connecting the supraglacial and englacial drainage networks and resulting in smaller-sized ponds (Miles et al., 2017b). However, as noted above, ponds are generally more likely to form in areas with lower surface velocities. Ponds may also drain by preferentially exploiting inherited structural weaknesses such as (sediment-filled) crevasse traces, open crevasses, and englacial conduits that have been forced closed by longitudinal compression, allowing drainage by hydrofracture (the penetration of a water-filled crevasse through an ice mass assisted by the additional pressure of the water at the crevasse tip) (Benn et al., 2017, 2012, 2009; Gulley and Benn, 2007; Miles et al., 2017b). Alternatively, perched ponds may drain by overspilling, when a channel is melted into the downstream end of a pond. If, during drainage, such a channel incises faster than the pond lowers then unstable and potentially catastrophic drainage can result (Liu et al., 2015; Raymond and Nolan, 2000). However, analyses on Lirung Glacier, Nepal Himalaya, provided strong evidence for continuous inefficient drainage of supraglacial ponds, likely into debris-choked englacial conduits (Miles et al., 2017a).

A periodic cycle of supraglacial pond expansion and drainage may occur until the pond becomes large enough to become permanently connected to the englacial system, and thus stabilise due to inputs of meltwater from streams and other ponds located farther upglacier (Benn et al., 2001; Miles et al., 2017a; Wessels et al., 2002). An abundant supply of meltwater from the

ice surface or the wider drainage system is evidenced by ponds with a high suspended-sediment concentration (Takeuchi et al., 2012). A seasonal pattern of supraglacial pond filling and drainage has been observed at seven glaciers in the Tien Shan, with 94% of observed ponds draining during the monsoon every year between 2013–2015 (Narama et al., 2017). Similar cycles were reported for five glaciers in Langtang Valley, Nepal Himalaya, where the maximum ponded area between 1999–2013 occurred early in the melt season, subsequently decreasing as ponds drained or froze (Miles et al., 2017b). Conversely, larger ponds have been observed to drain incompletely and separate into multiple smaller ponds, subsequently refilling to re-form one large pond (Benn et al., 2001; Miles et al., 2017b; Wessels et al., 2002). Warmer spring temperatures have been noted to correlate with a greater number of drainage events later the same year, likely due to larger meltwater inputs earlier in the year triggering redevelopment of the subsurface drainage system (Liu et al., 2015).

Supraglacial ponds are responsible for a large proportion of debris-covered glacier ablation, absorbing heat up to 14 times more quickly than even the debris-covered area. In the Langtang Valley, Nepal, this accounted for 12.5% of catchment ice loss (E. S. Miles et al., 2018b). However, linked supraglacial pond chains have been suggested to provide only a small proportion of total glacier proglacial discharge (Irvine-Fynn et al., 2017; Miles et al., 2019), primarily storing meltwater and thus increasing the potential for enhanced ablation. Ponds release $\geq 50\%$ of their absorbed energy with the melt output from the pond, contributing to internal melting along supraglacial and englacial conduits (Miles et al., 2016; Sakai et al., 2000). This in turn may lead to englacial roof collapse and the formation of new ponds (Benn et al., 2012; Miles et al., 2017a; Sakai et al., 2000), resulting in a net glacier-wide increase in ablation. The growing presence of ponds has been described as the clearest indicator of the influence of climate change on debris-covered glaciers (Salerno et al., 2012).

2.1.3 Meltwater transport

Supraglacial streams (Figure 6) on High Mountain Asian debris-covered glaciers vary widely in prevalence, size, and length. To exist and persist, a supply catchment is required (Benn et al., 2017; Gulley et al., 2009a) and the rate of stream incision, driven by thermal erosion, must outpace the rate of surface lowering (Marston, 1983). Such conditions may be promoted beneath thicker debris that suppresses surface ablation in the lower ablation area (Benn et al., 2017), yet observations of streams in this region are rare, likely due to the hummocky topography both limiting the size of supraglacial catchments (Fyffe et al., 2019b) and preventing any streams that do form from persisting for long distances (Benn et al., 2017). Farther upglacier, often under conditions of strong longitudinal extension associated with ice falls, open crevasses are common and also suppress supraglacial stream development (Benn et al., 2017). Most supraglacial streams have therefore been observed in the upper to mid-ablation area (Figure 6A-D) (Gulley et al., 2009a; Miles et al., 2019), downglacier of crevasse fields but still upglacier of the pronounced hummocky topography and thick debris layer (Section 2.1.1).

A perennial supraglacial stream (Figure 6A-D) has been present in the upper ablation area of Khumbu Glacier, Nepal Himalaya, for over 14 years (Gulley et al., 2009a; Miles et al., 2019). This stream and its smaller tributaries originate just downglacier of the Khumbu icefall, where the mean

longitudinal surface gradient decreases dramatically (Figure 1), from $\sim 23^\circ$ down the icefall to $\sim 3^\circ$ just below the clean-ice region (estimated from Shean (2017) over one km segments of the glacier's central flowline). The low surface gradient of the ablation area results in this channel having a high sinuosity (Miles et al., 2019). As streams transfer meltwater downglacier, they can incise effectively into the glacier surface (Figure 6B and C); one channel had melted 5–10 m deep by the time it reached the lower ablation area (Gulley et al., 2009a; Iwata et al., 1980). Such incision is evident where the channel sides and surrounding glacier surface have ablated more slowly than the channel itself, leaving walls of horizontally notched ice showing former, less incised channel elevations (Figure 6C). Supraglacial streams may drain into debris-covered glaciers through crevasses or moulins (Gulley et al., 2009a; Iwata et al., 1980), or through channel 'cut-and-closure' (see Section 3.1) (Gulley et al., 2009a; Jarosch and Gudmundsson, 2012). Relict channels abandoned by continued incision can often be exposed on the surface as a result of spatially variable surface lowering (Figure 6D).




Figure 6 – Examples of supraglacial streams on Khumbu Glacier, Nepal Himalaya, in: A-C) the upper ablation area, incised into the ice beneath the debris layer. Blue arrows indicate water flow direction; yellow arrows indicate abandoned/relict channels. The supraglacial stream in A) is extensive, sinuous, and very well developed, transporting large volumes of meltwater efficiently. B) and C) are upstream of A) (white and black star, respectively): B) shows a relict, debris-filled meander bend which has been superseded by a more direct channel; C) shows multiple levels of stream incision (grooves indicated by red dashed line, ~ 1 m high); D) the mid-ablation area, where the same incised channel becomes englacial through cut-and-closure after several hundred metres of progressive downcutting, visible from the multiple relict levels (channel drop in the image is ~ 10 m); E) and F) the lower ablation area. The channel in E) is a short stretch between a supraglacial pond and a shallow moulin, flowing over the debris layer. The stream in F) flows into a breach in the lateral moraine to form the proglacial stream; here, it has eroded into the sand-like sediment across a basin that seasonally floods. Images in A, E, and F were taken during fluorescein dye

tracing experiments (Miles et al., 2019). Image credit for A: Duncan Quincey; B, C, and E: Katie Miles; D: Evan Miles; and F: Bryn Hubbard.

Supraglacial streams can undergo rapid pathway changes. Figure 6B shows a debris-filled section of channel, abandoned as meltwater progressively took a more direct route, leaving a central island of protruding ice. This process may have been similar to the formation of an ox-bow lake in an abandoned terrestrial river meander. However, the abandoned channel section may be reactivated during times of high flow, evidenced by the presence of thick, evenly spread debris deposits in Figure 6B. Farther downglacier, where supraglacial stream observations are rarer, pathway changes have also been witnessed on short timescales (Miles et al., 2019). In Figure 6E, the stream flows into a shallow moulin, yet within 10 days this moulin had collapsed and been abandoned, with the stream routing into a new moulin just upstream. Moulin collapse has been attributed to the highly spatially variable surface lowering and ablation rates on debris-covered glaciers (Miles et al., 2019), while the short timescale indicates that the new moulin exploited an existing weakness in the ice.

2.2 Supraglacial knowledge gaps

Predictions of future mass balance regimes on High Mountain Asian debris-covered glaciers are uncertain. Surface lowering is leading to both an overall increase in debris thickness (Gibson et al., 2017a) and an upglacier emergence of a thin supraglacial debris layer. These processes will likely further decrease albedo and increase surface meltwater production (thereby increasing surface lowering, potentially leading to a positive cycle until debris thickens sufficiently to insulate the surface) (Kirkbride and Warren, 1999; Stokes et al., 2007). Measuring meltwater production is crucial, but difficult beneath (thin) debris layers, and often impossible where access to the ice-debris interface is not feasible. More broadly, the future evolution of debris-covered glacier surface geometry remains inadequately addressed, for example, whether meltwater will primarily be transported rapidly off the glacier in channels or stored within large systems of linked supraglacial ponds, thereby buffering diurnal proglacial discharge.

On a finer scale, a detailed process understanding of meltwater storage and transport through supraglacial ponds and pond systems is lacking, particularly of water circulation within, between, and out of ponds. While often just one discrete, channelised output is visible, water has also been observed to seep beneath the debris layer and emerge in unexpected locations (Miles et al., 2019). There has been little focus on how these links between ponds will change as ponds expand and eventually coalesce. Volumetric measurements of supraglacial ponds are scarce, rendering it difficult to accurately calculate how much meltwater is being stored on the glacier surface. Additionally, little attention has been paid to the effect of debris (heated by solar radiation) falling into a pond on the pond temperature and thus its subaqueous melt rate.

Understanding of the various pathways and rates of meltwater transport across a debris-covered glacier surface would benefit from additional focused research. For example, supraglacial streams are commonly difficult to discern in debris-covered regions of the glacier surface; this is particularly true for smaller surface streams and diffuse flows, which are less easily located and consequently remain largely unreported. On a smaller scale, the occurrence of some ice ablation

beneath even a thick debris layer implies that during much of the ablation season, water must exist between the ice surface and the debris layer (McCarthy et al., 2017), likely as a thin but variable film. However, the planform structure and meltwater transport of any such drainage network remain unknown, although transport must subsequently occur as a saturated surface layer or - initially at least - as small, inefficient rivulets.

Water storage within and below the supraglacial debris layer is likely but unexplored. Such storage would introduce temporary delays in the transport of meltwater through the system, thus affecting meltwater hydrochemistry (Tranter et al., 2002, 1993), the development of other parts of the drainage network, and proglacial discharge. However, despite its importance in contrasting with standard models of supraglacial hydrology based on research at clean-ice glaciers, small-scale meltwater storage delays remain unknown, which at least partly results from the difficulty involved in gaining access to the ice-debris interface beneath thick surface debris. Similar issues are present for the hydrology of snowpacks overlying thick debris; the extent that the snowpack delays runoff and how much snowmelt enters the hydrological system are similarly unaddressed.

3. Englacial hydrology

3.1 Englacial zone

Exceptionally, englacial conduits at High Mountain Asian debris-covered glaciers have been at least as well explored by glaciologists as at clean-ice glaciers. Such exploration has been carried out primarily in the Nepal Himalaya, including at Khumbu Glacier (Gulley et al., 2009a), Ngozumpa Glacier (Benn et al., 2017, 2009; Gulley and Benn, 2007), Ama Dablam and Lhotse Glaciers (Gulley and Benn, 2007), as well as several debris-covered glaciers in the Tien Shan (Narama et al., 2017). Largely on the basis of such studies, Gulley et al. (2009) proposed three formation mechanisms for englacial conduits within debris-covered glaciers:

- I. Cut-and-closure type conduits appear to be particularly prevalent within High Mountain Asian debris-covered glaciers, relative to clean-ice counterparts. Since the process requires more rapid supraglacial channel incision than surface ablation, this prevalence could result from the presence of cold surface ice and/or surface debris, both impeding general surface lowering. Under such conditions, incision will continue to the hydrologic base-level of the glacier (Section 2.1.2), with englacial conduits forming by supraglacial channel closure from snow infill and, in some cases, by ice creep (Gulley et al., 2009b, 2009a). These conduits may be repeatedly abandoned and reactivated as water supply varies through the year. However, such conduits rarely close completely due to their shallow depth below the surface, and may contain sediment that provides lines of secondary permeability by which the conduit may subsequently be reactivated (Benn et al., 2009; Gulley et al., 2009a; Gulley and Benn, 2007). Cut-and-closure conduits have been reported on Khumbu (Gulley et al., 2009a) and Ngozumpa Glaciers (Thompson et al., 2012).
- II. Meltwater may aggregate to form englacial conduits by exploiting lines/planes of secondary permeability; for example, those left by relict cut-and-closure conduits or debris-filled and/or compressed former surface crevasses (Benn et al., 2012; Gulley et

al., 2009b; Gulley and Benn, 2007; E. S. Miles et al., 2018a). Along these low-permeability zones, discharge through the icy matrix leads to the development of enlarging lines of preferential flow due to viscous heat dissipation, eventually forming an englacial conduit (Benn et al., 2012).

III. Englacial conduits may also form by hydrofracturing (Benn et al., 2012, 2009; Gulley et al., 2009b), though this process is generally restricted to upper, debris-free areas where surface runoff can enter open crevasses (Benn et al., 2012). In the lower ablation area, low surface gradients, low strain, and compression reduce the capacity for crevassing. Conduit formation by hydrofracturing has been invoked in association with longitudinal crevasses on Khumbu Glacier (Benn et al., 2012, 2009), promoted by the combined effect of transverse stresses and high water pressure at the base of supraglacial lakes. Multiple stages of hydrofracture, followed by conduit closure through freeze-on, were interpreted from a series of successively lower niches eroded into pond walls (Benn et al., 2009).

If a stream exploits a crevasse for sufficient time, it forms a moulin, as on clean-ice glaciers. Although such instances are rare, steep-gradient moulins have been observed in the upper ablation area of some High Mountain Asian debris-covered glaciers (e.g. Southern Inylchek Glacier, Tien Shan and Baltoro Glacier, Pakistan Karakoram (Narama et al., 2017; Quincey et al., 2009)), and a shallow-gradient moulin reported in the lower ablation area of Khumbu Glacier (Figure 6E) (Miles et al., 2019). Indeed, explored englacial conduits, such as on Khumbu and Ngozumpa Glaciers, also had shallow gradients (Benn et al., 2017; Gulley et al., 2009a; Gulley and Benn, 2007), suggesting predominant formation in these instances by cut-and-closure rather than crevasse exploitation.


Englacial conduits have been observed at multiple elevations within High Mountain Asian debris-covered glaciers, often showing numerous levels of incision hypothesised to result from sequential supraglacial pond drainage events as the base-level has moved (Gulley et al., 2009a; Gulley and Benn, 2007). According to this model, each conduit can have varying local base-levels through time (Section 2.1.2), with elevations ultimately limited by the glacier's contemporary base-level, determined by the height at which water leaves the glacier (Gulley et al., 2009a; Gulley and Benn, 2007). Furthermore, as the surface gradient of the ablation area of debris-covered glaciers is typically very low, the hydraulic gradient (Shreve, 1972) is correspondingly low, encouraging meandering and the formation of sinuous englacial conduits (Miles et al., 2019), as observed on Khumbu and Ngozumpa Glaciers (Benn et al., 2017; Gulley and Benn, 2007).

Longer-distance water transport has been inferred to occur through perennial sub-marginal conduits, likely formed by cut-and-closure, located along the edge of debris-covered glaciers (Benn et al., 2017; Thompson et al., 2016). Such marginal features provide longer-distance and more hydraulically efficient pathways than conduits within the central glacier, due to the frequent presence of infilled crevasse traces that can be exploited by water flowing at the margins (Gulley and Benn, 2007). Centrally located englacial conduits may become re-exposed due to lowering of the surrounding surface, routing water back to the surface (Figure 7) (Miles et al.,

2019). This process may make these conduits more discontinuous, particularly when combined with the commonly hummocky topography (Miles et al., 2017a).

Englacial systems have been observed at shallow depths below the surfaces of High Mountain Asian debris-covered glaciers. These typically consist of short conduits (channelised, distributed or a combination), englacial reservoirs, and/or shallow moulines, primarily linking supraglacial ponds (Miles et al., 2017a, 2019; Narama et al., 2017). Such linked supraglacial-englacial systems may be created and/or maintained by supraglacial pond drainage into englacial conduits (Gulley and Benn, 2007; Narama et al., 2017). Narama et al. (2017) found that the seasonal drainage cycle of supraglacial ponds on seven Tien Shan glaciers was characterised by a connection to an established englacial drainage system later in the summer; 94% of ponds drained and connected during all three years studied. Englacial conduits may thus play an important role in the life cycles of perched ponds (Benn et al., 2017; Miles et al., 2017a).

Deeper englacial drainage networks can vary in efficiency in response to numerous factors, including supraglacial pond drainage events. On Dokriani Glacier, Garhwal Himalaya, englacial conduits were inferred to be efficient and active through the entire melt season, with proglacial discharge proportional to supraglacial water production (Hasnain and Thayyen, 1994). Conversely, on Khumbu Glacier, a channelised but inefficient englacial system was inferred in the pre-monsoon season (Miles et al., 2019). This system did not link to the supraglacial pond chain, but was routed to the surface close to the terminus, suggesting that deep englacial to shallow-englacial-supraglacial links are also possible. While this inefficient englacial system was characterised by slow transport velocities, previous observations of faster transit through Khumbu Glacier during the drainage of a tributary glacier's supraglacial pond implies that the system can adapt rapidly to greater meltwater inputs (E. S. Miles et al., 2018a; Miles et al., 2019).



*Figure 7 – A relict englacial conduit (~10 m in height) in the centre of an ice cliff on Khumbu Glacier, Nepal Himalaya, exposed after a drainage event of the associated supraglacial pond, viewed: **A)** from upglacier, and **B)** from downglacier. On the downglacier side, tens of metres of surface lowering has occurred and the previously englacial conduit is now visible from the surface, meandering and incising for ~200 m farther downglacier before flowing into a pond. Image credit for A: Evan Miles; and B: Katie Miles.*

The efficiency of englacial meltwater transport has also been noted to change through the melt season at High Mountain Asian debris-covered glaciers. The influx of large volumes of monsoon precipitation during the summer months may result in the reopening of englacial (and subglacial) conduits, giving potential for considerable englacial ablation (Benn et al., 2012); for a surface pond of 500 m², sufficient energy to melt ~2,600 m³ of temperate ice is released over a single monsoon season (Miles et al., 2016). This additional meltwater ultimately leads to conduit erosion (Miles et al., 2017b; Sakai et al., 2000), which may be further enhanced by pond drainage events, as the warmer drained water (Section 2.1.2) conveys large amounts of energy, adding further to total glacier mass loss (Benn et al., 2012; Miles et al., 2016; Sakai et al., 2000; Thompson et al., 2016).

For englacial conduits located near the surface, rapid expansion can result in conduit collapse if the ceiling is not sufficiently supported. A supraglacial, possibly relict, channel formed from englacial conduit collapse exposes new bare ice faces, including ice cliffs, which may then contribute to more rapid lowering of the glacier surface (Section 2.1.1) (Benn et al., 2017; Kraaijenbrink et al., 2016b; Miles et al., 2016; Sakai et al., 2000; Thompson et al., 2016, 2012). Ablation rates and surface subsidence can be further enhanced if the new depression becomes flooded by that increased meltwater production, supplemented by upglacier inputs, providing new depressions for supraglacial ponds to form or expand and coalesce (Section 2.1.2) (Benn et al., 2012, 2001; Kirkbride, 1993; Kraaijenbrink et al., 2016b; Miles et al., 2017a; Sakai et al., 2000; Thompson et al., 2012).

Meltwater may be stored englacially within debris-covered glaciers, ranging from small, shallow englacial reservoirs (Miles et al., 2019) to deeper and potentially larger reservoirs. The latter type has been inferred, for example, for glaciers feeding the Hunza river system, Central Karakoram, at the start of the melt season due to a notable lag between the initiation of glacier ablation and higher discharges observed downstream (Hewitt et al., 1989). Similarly, the initiation of an outburst flood at Lhotse Glacier was partly attributed to the release of meltwater stored within englacial conduits that became over-pressurised from greater meltwater production and transit during the transitional pre-monsoon season (Rounce et al., 2017). Other inferences have been made from supraglacial pond water-level measurements, such as at Imja Tsho, Nepal Himalaya, where the post-melt season lake level was constant despite lower air temperatures and lower precipitation, which would both serve to reduce meltwater production. This situation was explained by recharge from englacially and subglacially stored water being progressively released over time (Thakuri et al., 2016).

3.2 Englacial knowledge gaps

Despite relatively extensive englacial glacioclimatological exploration, numerous gaps remain in our knowledge of the englacial drainage of High Mountain Asian debris-covered glaciers. For example, as at clean-ice glaciers, the thermal regime of the glacier exerts a significant control on the location and formation of an englacial drainage system, yet the thermal regime is unknown for almost all High Mountain Asian debris-covered glaciers. A recent study suggested that the lower area of Khumbu Glacier may primarily comprise temperate ice (K. E. Miles et al., 2018) allowing the existence of a deep englacial drainage system (Miles et al., 2019). However, this research was

confined to a single glacier and its representativeness for other debris-covered glaciers in High Mountain Asia remains unknown.

Knowledge of the influence of supraglacial debris on englacial (and subglacial) drainage systems is incomplete. On Miage Glacier, the upglacier ice, which is cleaner and covered with a thin supraglacial debris layer, was shown to produce an efficient subsurface drainage system to the terminus from early in the melt season. In contrast, the heavily debris-covered lower ablation area restricted the development of supraglacial drainage, leading to an inefficient subsurface system that ultimately flowed into the efficient system driven from upglacier (Fyffe et al., 2019b). While there are similarities between the drainage system of Miage and the few High Mountain Asian debris-covered glaciers studied, the generally thicker debris layer and much greater prevalence of supraglacial ponds towards the terminus of the latter will additionally influence the hydrological system of such glaciers – an influence that remains unexplored.

Improved understanding is required of the links between the englacial system and other hydrological domains, such as supraglacial-to-englacial transitions (through cut-and-closure conduits, weaknesses in the ice, and supraglacial pond drainages). Research into the shallow englacial system is needed, including determining how much of a distinction there is between shallow englacial and supraglacial systems, considering the rapidly changing surface topography that is typical of High Mountain Asian debris-covered glaciers. Finally, the potential for englacial meltwater storage has received very little attention.

4. Subglacial hydrology

4.1 Subglacial zone

Knowledge of subglacial drainage at High Mountain Asian debris-covered glaciers is limited, although some evidence at least points to the existence of such systems. For example, glaciological investigations indicated that the proglacial stream of a retreating tributary of Khumbu Glacier reached Khumbu's bed (Benn, pers. comm., 2018). This conduit was assumed to follow the bed for some distance downglacier, similar to the perennial sub-marginal conduits present at the edge of the neighbouring Ngozumpa Glacier (Benn et al., 2017; Miles et al., 2019; Thompson et al., 2016). However, this water did not persist subglacially, and instead was documented to exit the glacier supraglacially. This upward routing of water likely occurs due to the commonly high hydrological base-level of such glaciers, possibly following the glacier's cold-temperate transition surface (K. E. Miles et al., 2018; Miles et al., 2019).

Beyond the studies outlined above, all other information relating to the subglacial drainage of High Mountain Asia debris-covered glaciers is inferred. For example, the presence of meltwater at the bed has been inferred from surface velocity records from remote sensing (e.g. Quincey et al., 2009) or field-based GPS (e.g. Tsutaki et al., 2019), using inferences similar to those for clean-ice glaciers. Relatively rapid surface velocities in the central areas of glaciers have been recorded during summer months, when melting and rainfall delivery are greatest (Figure 8). Such velocity increases have been interpreted as indicative of basal motion lubricated by subglacial drainage (Benn et al., 2017; Copland et al., 2009; Kääb, 2005; Kodama and Mae, 1976; Kraaijenbrink et al.,

2016a; Kumar and Dobhal, 1997; Mayer et al., 2006; Quincey et al., 2009). Similar remote sensing studies of surging debris-covered glaciers, particularly in the Karakoram, have inferred the presence of subglacial water that enables rapid surface velocities during surge phases (Copland et al., 2009; Quincey et al., 2011; Steiner et al., 2018a), such as the maximum velocity of $> 250 \text{ m a}^{-1}$ reported at South Skamri Glacier, Pakistan Karakoram (Copland et al., 2009).

The existence of channelised subglacial drainage has been inferred from the presence of proglacial outlet streams at the terminus of debris-covered glaciers. During the melt season, these channels discharge large volumes of heavily debris-laden water, implying sediment entrainment during transport along the bed (Quincey et al., 2009). This transport pathway has also been inferred from comparisons of supraglacial with proglacial solute concentrations on Lirung Glacier, where high proglacial Ca^{2+} and SO_4^{2-} concentrations indicated prolonged contact with reactive debris, inferred to occur during subglacial drainage (Bhatt et al., 2007). Similarly, a perennially active subglacial system on Dokriani Glacier was inferred to be connected with the englacial system on the basis of proglacial electrical conductivity measurements (Hasnain and Thayyen, 1994).

Variations in subglacial system efficiency have been inferred from studies focusing on proglacial streams. For example, the increasing efficiency and interconnection of the subglacial system of Gangotri Glacier, Garhwal Himalaya, was inferred from an increase in the net flux and size of subglacially eroded suspended particles through the melt season (Haritashya et al., 2010). On the same glacier, dye tracing experiments showed that the channelised subglacial drainage system became progressively more efficient with greater meltwater inputs through the melt season (Pottakkal et al., 2014). Dye tracing experiments have also demonstrated a transition from distributed to channelised subglacial drainage through the melt season, for example at both Dokriani Glacier and Hailuoguo Glacier, Mt. Gongga, Tibet (Hasnain et al., 2001; Liu et al., 2018). On a diurnal scale, Kumar et al. (2009) found that the total ion concentration of proglacial meltwater at Gangotri Glacier increased from the afternoon onwards, interpreted as an enhanced subglacial component due to the englacial system developing through the day and transporting a greater proportion of supraglacial meltwater to the solute-rich glacier bed. Finally, substantial subglacial meltwater storage at debris-covered Lirung Glacier was inferred from its lower diurnal discharge variability relative to nearby debris-free Khimsung Glacier, Nepal Himalaya (Wilson et al., 2016).




Figure 8 – Surface velocity maps of Lirung Glacier, Nepal Himalaya, during summer (left) and winter (right), with three transverse velocity profiles (A–C) at the locations marked. From Kraaijenbrink et al. (2016b).

4.2 Subglacial knowledge gaps

Very little is known about the subglacial drainage of High Mountain Asian debris-covered glaciers, largely due to the difficulty in accessing these systems. Furthermore, many debris-covered glaciers in High Mountain Asia terminate in lakes (Section 5.1.1), which increases the likelihood of some form of subglacial drainage system but reduces the likelihood of that system being channelised. Such lakes also severely hamper direct access to any outflow streams that might be present. Assuming the existence of such conduits, it is entirely unknown whether subglacial networks flow directly into proglacial ponds at the bed, are routed to the surface upglacier and flow in supraglacially (similar to the pathway of some englacial drainage at Ngozumpa Glacier (Benn et al., 2017)), or are partially or wholly lost to groundwater. Additionally, the existence of base-level englacial streams and a perched water table are highly likely to complicate the detection of, and distinction between, englacial and subglacial systems, at least approaching the terminus. For example, towards the terminus of Khumbu Glacier, it has been inferred that the high local base-level results in the uprouting of the subglacial/deep englacial drainage system to the surface, yet, since the ice here is temperate, some meltwater would nonetheless be expected at the bed (K. E. Miles et al., 2018; Miles et al., 2019). However, basal ice temperatures and conditions for almost all other High Mountain Asian debris-covered glaciers are entirely unknown.

Transitions between the englacial and subglacial system are important to understand, as are discovering and tracking lost meltwater components – lost potentially to groundwater, to short- or long-term storage within the glacier, or to evaporation from the terminal moraine. If

extensive subglacial drainage environments are discovered, the influence of the supraglacial debris cover on those systems should also be investigated.

5. Proglacial hydrology

5.1 Proglacial zone

5.1.1 Proglacial lakes

One of the most distinctive characteristics of the proglacial zone of High Mountain Asian debris-covered glaciers is the frequent presence of a proglacial lake (Figure 9), which are far more common than at equivalent clean-ice glaciers. These lakes form by a continuation of the processes of glacier thinning and supraglacial pond growth (Section 2.1.2) facilitated by the deposition of sufficient debris by debris-covered glaciers to create high, arcuate terminal moraines. Here, perched supraglacial ponds expand both downwards, eventually cutting to base-level, and laterally, often eventually coalescing to produce one large lake above and over the terminus (Basnett et al., 2013; Kattelmann, 2003; Mertes et al., 2016; Röhl, 2008; Watanabe et al., 2009). Although less common, base-level lakes that penetrate the full glacier thickness can form farther upglacier and expand downglacier through stagnant terminus ice, for example Imja Tsho on Imja-Lhotse Shar Glacier, Nepal Himalaya (Figure 9) (Watanabe et al., 2009). The exact location of such a proglacial lake may be determined by the location of shallow englacial conduits that provide pre-existing lines of weakness as the perched ponds grow (Benn et al., 2017; Thompson et al., 2012). Proglacial lakes will therefore determine the hydrological base-level of the glacier, and are often dammed by the terminal moraine (Thompson et al., 2012).




Figure 9 – Proglacial lake (Imja Tsho) with a frozen and snow-covered surface at Imja-Lhotse Shar Glacier, Nepal Himalaya. A) full length of Imja Tsho (~2.7 km in October 2018), looking upstream towards the calving front of Imja-Lhotse Shar Glacier. B) detached (stagnant) glacier ice that dams the lake. The black star and arrow in B) show the location and direction A) was taken in. Image credits: Katie Miles.

The formation of moraine-dammed proglacial lakes represents a final stage in the surface lowering and overall mass loss of debris-covered glaciers. Benn et al. (2012) defined three stages in the development of debris-covered glaciers: in regime one, all parts of the glacier are

dynamically active; in regime two, surface lowering has begun and ice velocities decrease; in regime three, glaciers are stagnant and rapid recession may occur. The formation of a base-level lake indicates that a glacier has entered this third regime, and rapid recession may then occur through further expansion of that proglacial lake (Benn et al., 2012). A growing number of proglacial lakes of increasing size have been observed in recent decades across the Hindu Kush Himalaya (Gardelle et al., 2011; Haritashya et al., 2018b; Nie et al., 2017; Thompson et al., 2012). The pattern of proglacial lake formation varies across the region, with proglacial lake area in the western Himalaya decreasing 30–50% from 1990–2009 compared to an increase of 20–65% towards the east, where proglacial lakes are already more prevalent (Gardelle et al., 2011; Maharjan et al., 2018). This pattern at least partly results from greater glacier recession in the west over this period (Gardelle et al., 2011).

Proglacial lakes expand through similar mechanisms to supraglacial ponds (i.e. subaqueous melting and subaerial ice face melting; Section 2.1.2) until they are limited by substrate. Lake expansion therefore enhances glacial mass loss and meltwater production while the lake is underlain or dammed by ice (Carrivick and Tweed, 2013; Röhl, 2008). Once calving is triggered, it becomes the dominant method of subsequent lake growth (Röhl, 2008; Thompson et al., 2012). Calving into a proglacial lake progresses from notch development and roof collapse to large-scale, full-height slab calving enabled by the lake deepening to the glacier bed (Kirkbride and Warren, 1997; Thompson et al., 2012). The water depth may then be sufficient to trigger extending flow in the now-unsupported ice cliff, increasing flow velocities and weakening the ice through crevasse formation and dynamically induced thinning (King et al., 2019; Kirkbride and Warren, 1999; Thompson et al., 2012; Tsutaki et al., 2019). This can result in rapid and potentially unstable calving, substantially increasing glacier mass loss, as has been observed during several kilometres of such retreat at Tasman Glacier, New Zealand (Kirkbride and Warren, 1999) and modelled for lake- and land-terminating glaciers in the Bhutan Himalaya (Tsutaki et al., 2019). Upglacier expansion of the proglacial lake (Watanabe et al., 2009) may have implications for the glacier's drainage system, such as by earlier interruption of meltwater routing (Carrivick and Tweed, 2013).

Very large proglacial lakes can alter a glacier's microclimate due to a lake's lower albedo and higher thermal heat capacity relative to the surrounding ice and soil, thereby producing locally cooler summer air temperatures and warmer autumn temperatures (Carrivick and Tweed, 2013). This can slow local summer ice ablation and consequently reduce the amount of meltwater being produced and transported through the glacier, with implications for the development of englacial and subglacial drainage systems. If a moraine-dammed proglacial lake is present then the overwhelming majority of water transported through a debris-covered glacier is likely to pass through that lake (Benn et al., 2017). This has implications for drainage through the glacier and for the potential occurrence of glacial lake outburst floods (GLOFs).

5.1.2 Proglacial streams

Proglacial runoff from debris-covered glaciers can form a significant proportion of the discharge of large rivers downstream, particularly in High Mountain Asia: the Indus, Dudh Koshi, Ganges, and Brahmaputra rivers all stem from glacial meltwaters (Pritchard, 2019; Ragettli et al., 2015; Wilson et al., 2016). In particular, glacial runoff buffers both seasonal (Bolch et al., 2019; Pritchard, 2019)

and annual (Pohl et al., 2017) water shortages. Loss of glacier volume due to longer, warmer melt seasons and decreased snow accumulation could result in periods of much reduced water availability, greatly influencing downstream communities and ecology (Bolch et al., 2019; Pohl et al., 2017; Pritchard, 2019).

Proglacial discharge measurements, estimates, and models have been run across High Mountain Asia, such as on individual glaciers in Nepal (Braun et al., 1993; Fujita and Sakai, 2014; Ragettli et al., 2015; Rana et al., 1997; Savéan et al., 2015; Soncini et al., 2016; Tangborn and Rana, 2000), Tibet (Kehrwald et al., 2008), the Tien Shan (Chen and Ding, 2009; Han et al., 2010; Sorg et al., 2012), India (Hasnain, 1999, 1996; Khan et al., 2017; Singh et al., 2005, 1995; Singh and Bengtsson, 2004; Thayyen and Gergan, 2010), and for multiple catchments and entire regions (Winiger et al., 2005). However, such records are relatively short: of the studies listed above, five measured discharge for a year or less; three have 2–3 years of measurements; and only one has 6 years of measurements. The others use modelling to obtain estimates of proglacial discharge.

The presence of surface debris can have a notable effect on a glacier's proglacial discharge, resulting in a proglacial hydrograph that is different from that of a clean-ice glacier. While no such comparison has been made for a High Mountain Asian debris-covered glacier, an example is shown from the debris-covered Dome Glacier, Canadian Rockies (Figure 10) (Mattson, 2000). Here, discharge was muted both diurnally and through the ablation season compared to the neighbouring clean-ice Athabasca Glacier (Figure 10); variation in annual discharge volume from Dome Glacier between the two years was 1%, compared to 24% from Athabasca Glacier. This is due partly to the suppression of surface melt by a debris cover (Section 2.1.1), and partly to the lags that are induced as a result of the debris layer – the additional time to conduct heat through the debris and the warmer local air temperatures due to the warming debris introduces a delay. Thus, peak melt can occur up to several hours after the maximum radiation receipt at the debris surface (Carenzo et al., 2016; Conway and Rasmussen, 2000; Evatt et al., 2015), and an exceptional case has been recorded as being up to 24 hours later for debris layers > 0.85 m thick (Fyffe et al., 2014). This lag in diurnal peak melt is thus reflected in the timing of the highest stream flow, producing a later and less pronounced peak in the diurnal pattern of a debris-covered glacier's proglacial stream (Fyffe et al., 2019a, 2014).




Figure 10 – Hydrographs of proglacial discharge of the clean-ice Athabasca Glacier and the adjacent debris-covered Dome Glacier, Canadian Rockies, over the ablation months of July and August 1994 and 1995. Redrawn from Mattson (2000).

Lags in proglacial discharge from debris-covered glaciers may also be caused by the temporary storage of water within the surface debris layer, for example, during rainfall events. This may influence subglacial and proglacial discharge by delaying and buffering water transfer at the surface, potentially affecting basal water pressures and minimising peaks in proglacial discharge (Brock et al., 2010). In the Himalaya, monsoon precipitation is thought to exert a significant control on proglacial discharge hydrographs at high rainfall intensities. For example, Thayyen et al. (2005) suggested such an intensity was $> \sim 20 \text{ mm d}^{-1}$, which occurred on 20% of rainfall days during four years of monsoon measurements on Dokriani Glacier. Early in the melt season, meltwater is also stored within the snowpack of debris-covered glaciers, providing a further delay in the transport of meltwater from the surface into the subsurface drainage system (Singh et al., 2006b). In the last two decades the amount of snowfall accumulation has decreased across the Himalaya, and is projected to decrease a further 20–40% by 2100 (Salerno et al., 2015; Smith and Bookhagen, 2018; Viste and Sorteberg, 2015); this buffer will thus be further reduced, influencing the future proglacial hydrograph pattern of debris-covered glaciers.

Groundwater stored within high-elevation glacial catchments has been inferred to interact with proglacial (and subglacial) stream networks, affecting their discharge patterns due to additional water storage and subsequent release (Gremaud et al., 2009; Smart, 1996, 1988). For example, a ~ 45 day lag between precipitation and discharge was observed for 12 glacierised and non-glacierised Himalayan catchments, indicating storage of up to two-thirds of the river discharge in a groundwater aquifer system before the monsoon, greatly affecting the annual discharge pattern (Andermann et al., 2012c). This has similarly been shown by much reduced river

suspended sediment concentrations measured post-monsoon, having been diluted as groundwater begins to be released (Andermann et al., 2012b, 2012a). Comparable processes may occur beneath the glaciers themselves, for example, at Khumbu Glacier in the pre-monsoon season, where more meltwater entered the glacier's subsurface drainage system than exited the glacier at the terminus (Miles et al., 2019). Indeed, in the Jade Dragon Snow Mountain region of southwest China, 29% of glacier meltwater was calculated to be stored in a karst aquifer (Zeng et al., 2015). Groundwater sinks of subglacial meltwater can therefore comprise a significant portion of the total glacial output, potentially resulting in underestimation of glacial ablation.

A range of models has been used to predict future runoff from debris-covered glaciers using various future climatic scenarios for a single glacier basin (Ragettli et al., 2015; Singh et al., 2008, 2006a; Zhang et al., 2007) and multiple glacier basins (Immerzeel et al., 2012; Lowe and Collins, 2001) up to a regional scale (Rees and Collins, 2006; Shea and Immerzeel, 2016). Currently, a large proportion of debris-covered glaciers worldwide, particularly in the Himalaya, have negative mass balances (Bolch et al., 2012, 2011; Kääb et al., 2012; Scherler et al., 2011). A recently observed decline in Himalayan snowfall will contribute further to the decreasing mass of these glaciers by both reducing accumulation rates and exposing the glacier surface to atmospheric melting earlier in the melt season (Salerno et al., 2015). Glacier contributions to catchment discharge in many regions have been predicted to increase over the next few decades, but as the glaciers continue to shrink, peak water will be surpassed and this proportion will begin to reduce substantially due to the reduced volume of the remaining glaciers (Barnett et al., 2005; Bolch, 2017; Bolch et al., 2012; Huss, 2011; Huss and Hock, 2018; Lutz et al., 2014). Shea and Immerzeel (2016) estimated that most basins will have declining glacier contributions to streamflow by 2100, and water shortages may then be a concern for many populated areas in the Karakoram, while reduced peak flows may represent a greater concern in the eastern Himalaya.

5.2 Proglacial knowledge gaps

Few glacial discharge monitoring stations have been in place for longer than a decade in High Mountain Asia, leaving current and future discharge volumes unknown for most debris-covered glaciers. The volume and temporal variability of potential glacial meltwater losses to groundwater, and whether these re-join the glacial system (subglacially, proglacially, or further downstream), are also poorly understood.

Projections of future changes in proglacial hydrology are hampered by the absence of accurate predictions of the future geometric development of High Mountain Asian debris-covered glaciers. For example, if surface lowering remains the dominant response to climate warming, glaciers may melt entirely and/or form large proglacial lakes that then dominate mass loss processes. Conversely, the inverted mass balance regime could result in a separation of stagnant, heavily debris-covered lower glacier tongues from the upper, less debris-covered regions, potentially providing ideal conditions for a base-level lake to form in between, dammed by the detached debris-covered ice.

6. Future research themes

Based on the review above, we identify six hydrological research themes, including examples of appropriate techniques, that would contribute substantially to advancing our understanding of the hydrology of High Mountain Asian debris-covered glaciers.

I. Elucidating glacier-wide water balance

Given the importance of glaciers as a source of water in high mountain regions (Immerzeel et al., 2020), more robust quantification of water inputs into, and outputs from, the glacier system is paramount. Detailed and temporally and spatially extensive hydrological field measurements are required to better constrain numerical model parameterisations. Water inputs should be simulated and examined independently of glacier-fed river discharge, with attention to process parameterisation to facilitate improvements in efforts to close the water balance. Water storage is also an important component of the water balance, discussed further in research theme IV below.

The limited measurement to date of precipitation across High Mountain Asia, particularly in terms of partitioned snow and rainfall and synoptic and seasonal-to-annual variations in precipitation gradient and rainfall fraction, should be augmented by establishing a network of robust automatic weather stations over a range of catchments, surface types, and elevations. Glacier surface elevation change should be measured simultaneously by remote sensing and ground-based methods – for example, by ultrasonic rangiers – to calibrate and validate models of melt and mass balance. These approaches would also aid in determining the impact of anthropogenic black carbon aerosols and other light-absorbing impurities on albedo, supported by remote-sensing studies of surface characteristics. Precipitation gradients should be quantified further through dedicated accumulation measurements.

The retention of meltwater, for example by refreezing of meltwater within supraglacial debris, firn, crevasses, or the body of the glacier, requires better characterisation. Empirical data collected from snow pits and shallow ice cores would be sufficient to quantify such mass retention over short timescales, complemented by longer-term records derived from deeper coring or visual examination of layering present in borehole walls. In the accumulation area, these methods would provide the additional bonus of historical records of local accumulation.

The amount of water lost through evaporation and sublimation should be assessed through comprehensive studies of eddy covariance coupled with meteorological measurements. Future research should examine these processes not only from snow-covered areas, recently shown to be a key source of water loss (Stigter et al., 2018), but also over the accumulation and debris-covered ablation areas and the terminal and lower lateral moraines, which may equally contribute to evaporation and sublimation losses. Quantifying these moisture fluxes may be possible either by direct field measurement or by remote sensing for longer timescales.

Other research needs include quantifying losses to groundwater and better evaluating the role of debris in driving the observed hysteretic behaviour of downstream annual hydrographs. Hydrochemical and isotopic analyses may shed light on water sources and variations therein, while catchment-scale dye or gas tracing studies tied closely to continuous measurements of discharge at various locations on and beyond the glacier could help to define the volumes of water delivered to groundwater systems (and if so, the proportion that re-joins the proglacial stream farther downvalley).

II. Understanding hydrological processes influencing glacier mass balance

The efficiency of rainfall and meltwater routing from higher elevation locations should be evaluated due to its potential effect on glacier accumulation and mass balance by englacial melting. For example, heat fluxes driven by meltwater conveyance to the englacial and subglacial environments of debris-covered glaciers (i.e. cryo-hydrologic warming (Gilbert et al., 2020; Phillips et al., 2010)), could be explored using numerical models guided by field-based measurements of supraglacial water fluxes and temperatures, along with geophysical and/or borehole-based investigations of englacial temperature fields. Specific loci and timescales of meltwater routing, storage, and release should be determined. Englacial drainage pathways are of particular importance due to their strong association with the formation of supraglacial ice cliffs, which account for a disproportionate amount of surface melt at heavily debris-covered glaciers. Investigations should map current streams and monitor changes in surface topography and hydrology (for example, the collapse and surface exposure of shallow englacial systems) both remotely, using satellite images where streams are large enough, and in the field. The latter should be supplemented by dye tracing experiments to characterise the hydraulic properties of englacial systems.

There is a need for accurate knowledge of spatial variations in surface debris characteristics and thicknesses, and of meltwater located at the ice-debris interface, to improve models of surface vapour fluxes. Thus, meteorological stations are required to measure water content or relative humidity. Debris thickness maps and the existence of ponded and surface water could be constructed by refining algorithms from remotely sensed data (both thermal imagery and surface lowering) or on the basis of manual field measurements of ponds and high-frequency ground-penetrating radar to identify water present at the interface between the supraglacial debris layer and the underlying ice. Future investigations of supraglacial pond expansion rates should focus on wide-scale systematic field-based bathymetry, pond-sediment stratigraphic assessment, and measurements of pond water and basal sediment temperatures at multiple depths (particularly to assess vertical heat transfer from warm supraglacial pond water to the base of the pond), combined with the development of numerical models of heat transfer by such mechanisms.

III. Identifying the influence of drainage and meltwater storage on ice motion

Meltwater present at the bed or the terminus of debris-covered glaciers can affect the velocity of both land- and lake-terminating glaciers; a better understanding and inclusion of subglacial hydrological processes into models of glacier dynamics will improve future simulations of ice flow and glacier evolution. Within subglacial hydrological processes, better quantification is needed of

the inputs to the system (i.e. coupling meteorological data with melt modelling), the volume of water present at the bed (for example by monitoring subglacial water pressure in deep borehole arrays), and the volumes of water lost from the system (i.e. by calculating the glacier's water balance).

Ice motion should be separated into its constituent components (i.e. ice deformation and basal motion), with particular focus on measurements acquired during the melt season and on an individual glacier scale. Basal water pressure, and consequently glacier sliding, should be estimated through analysis of variations in glacier surface velocity obtained, for example through combining remote-sensing data with field-based GPS studies. The recently available and constantly growing archive of rapid-repeat, high-resolution optical and radar remotely sensed imagery will help future work to improve knowledge of seasonal velocities (e.g. Dehecq et al., 2019). Deeper ice velocities and strain can be recorded within boreholes, ideally extending to the glacier bed. Such boreholes can also allow measurements of the glacier thermal regime and bed substrate, while improved mapping of glacier bed topography across High Mountain Asia is necessary to constrain estimates of ice thicknesses. Finally, in order to assess the influence of calving from proglacial lakes, the above measurements should be collected in comparative studies of both lake- and land-terminating High Mountain Asian glaciers.

IV. Characterising seasonal changes in hydrology

Targeted research is needed to measure seasonal changes in hydrological storage components, particularly those that are specific to debris-covered glaciers. Improved understanding of storage components is needed to represent the drainage system of debris-covered glaciers appropriately in hydrological models. For example, seasonal changes in the area and volume of perched supraglacial ponds could be achieved at the glacier scale using rapid-repeat optical satellite imagery to maximise likelihood of observation and/or by using high-resolution synthetic aperture radar satellite data, which are insensitive to cloud cover. Detailed examination of the water content of the supraglacial debris layer (including the seasonal thaw dynamics of the debris layer, influencing its hydraulic transmissivity) can be made using soil moisture sensors installed at multiple depth intervals, while through-debris transmissivity and snowpack storage/release could be assessed by dye tracing experiments. These processes will aid better understanding of the role of debris, snow, and firn in transmitting meltwater to supraglacial streams and the subsurface drainage system, including seasonal storage and release from subsurface reservoirs.

Process-based understanding of seasonal hydrological changes could also be improved by detailed field-based studies. Glacier drainage systems respond dynamically to the seasonal production of meltwater; this is clear at clean-ice glaciers where snowline retreat stimulates the progressive upglacier transition from inefficient to efficient drainage. Research is needed at High Mountain Asian debris-covered glaciers to evaluate whether distinctive seasonal dynamics can be explained by additional storage components or specific melt-generation patterns. These phenomena can be addressed through dye tracing, glacioclimatology or bulk proglacial meltwater analysis. Such studies would also result in a better general understanding of the nature and form of englacial and subglacial drainage at High Mountain Asian debris-covered glaciers.

Finally, the seasonal structure and dynamics of debris-covered glacier hydrological systems should be understood in the context of projected future melt and discharge. An integrated effort to assess seasonal changes in debris-covered glacier hydrology should be coupled with melt season meteorological and ablation measurements, as well as development of a continuous discharge record through proglacial discharge monitoring stations.

V. Evaluating hydrological hazards

The growth in both number and size of supraglacial ponds is one of the clearest visual signs of debris-covered glacier decay. Future research should focus on predicting formation and growth of such ponds by combining glacier melt projections (e.g. Kraaijenbrink et al., 2017; Rounce et al., 2020) with modelled glacier bed overdeepenings (e.g. Linsbauer et al., 2016). Moraine-impounded sites (such as where base-level terminal lakes have been observed to develop) are more complex; investigations of the drainage capacity (evidence of free-drainage as opposed to impoundment), combined with remotely sensed observations of expanding and coalescing supraglacial pond chains, may provide a suitable starting point. Improved understanding of supraglacial pond expansion rates, discussed in research theme II, is also crucial, while accurately modelling the longevity of ice cliffs could be improved with high-resolution digital elevation models (obtained, for example, through Structure-from-Motion) coupled with simple numerical melt modelling.

Assessments of how ‘dangerous’ a lake is (potential for a catastrophic GLOF occurring) often disagree (e.g. Haritashya et al., 2018a; Maharjan et al., 2018; Rounce et al., 2016) and, while recent events such as the 2015 Gorkha earthquake suggest that the terminal moraines of glacial lakes may be more stable than hitherto considered, large-scale remote observations cannot assess internal or small-scale superficial damage caused by such events (Byers et al., 2017; Kargel et al., 2016). Such studies should be improved in terms of their sophistication, for example addressing a broader range of factors that contribute to the formation of a hazardous lake (e.g. Rounce et al., 2016), many of which may be site specific. Traditional magnitude-frequency relationships may no longer be relevant as the current state of mountain environments is non-stationary and beyond historic precedence. Therefore, alternative forms of event prediction are needed, such as site-specific hazard development depending on different event magnitudes.

Field-based measurements should be made on, and downstream of, individual proglacial lakes to determine potential hazards and the GLOF risk. Knowledge of moraine dam composition (including sediment type and the presence or absence of an ice core) and the existence of seepage or piping is needed, and could be addressed by radar, seismic studies, or drilling into moraines to characterise soil strength and composition. Flood hydrographs could be better constrained by geotechnical modelling to understand dam failure mechanisms. While predicting the timing of an outburst flood is nearly impossible, particularly those originating from englacial and subglacial sources, characterising subsurface drainage and routing and seasonal release of stored water may help to identify likely timing and locations of sudden outbursts (research theme IV). Cascading hydrological hazards, which may be triggered by very high-elevation and often hanging glaciers that are seldom studied, should also be considered. The thermal conditions and hydrology of these

glaciers should be investigated, for example, by surface ground-penetrating radar guided by borehole-based sensors, dye tracing and discharge monitoring.

VI. Predicting future hydrological changes over short and long timescales

Understanding the timescales over which debris-covered glaciers will lose mass, thereby influencing the amount of meltwater generated and subsequent hydrological processes, depends on developing a new generation of detailed glacier models that capture both the complex feedbacks between debris transport by ice and the processes affecting sub-debris ablation over timescales longer than a few decades (Rowan et al., 2015). Numerical model predictions need to integrate opposing processes on different scales, for example, encompassing both the glacier-scale 'debris-cover anomaly' (the observed, but still unexplained, debris-covered glacier mass loss rates that are similar to those of clean-ice glaciers (Brun et al., 2019; Gardelle et al., 2012; Pellicciotti et al., 2015)) and the smaller-scale insulating effect of the debris. Field and remote-sensing data relating to mass balance and ice flow processes are required at the correct scale and resolution for use in numerical models of glacier mass change, parameterised with sufficient process-based understanding to predict how these controls will evolve over time. The inclusion of these small-scale and complex processes within regional models (e.g. Kraaijenbrink et al., 2017) to improve the accuracy of large-scale mass-loss predictions should also be explored.

As debris-covered glaciers get smaller, primarily by surface lowering, the debris cover will thicken and increase insulation, reducing ablation over a potentially greater area of the terminus. Debris-covered glaciers are therefore already larger and likely to decline slower than equivalent clean-ice glaciers; as a result, the meltwater of clean-ice glaciers will temporarily provide a relatively larger component of the annual hydrological budget as they lose mass preferentially. Robust dynamic glacier models are therefore needed to predict changing hydrographs and contributions to downstream water supplies, particularly as peak water passes. Supraglacial ponds play an important role in modulating the proglacial hydrograph and, in the long-term, may provide a natural water supply reservoir during periods of drought. However, sedimentation rates within ponds, and therefore their likely longevity, should be quantified by *in situ* hydrological stations.

The acceleration of debris-covered glacier mass loss and decrease in glacial runoff as peak water passes are likely to lead to proglacial streams becoming proportionately more sediment-laden. This may be enhanced during the melt season, particularly in regions of High Mountain Asia affected by heavy monsoon rains, which can enhance supraglacial debris erosion. Furthermore, ice within larger debris-covered glaciers is older than in smaller glaciers and will thus contain a longer legacy of environmental contaminants (e.g. Hodson, 2014; Li et al., 2017). Ultimately, this may result in more pollutants being delivered via proglacial streams to water supplies, particularly during the melt season. Discharge and water quality should therefore be monitored with hydrological monitoring stations on proglacial streams across High Mountain Asia. Combined with modelling efforts and improved hydrological understanding, this will allow mitigation strategies to be planned for the vast downstream populations that depend on that meltwater.

944 7. Summary

945 In this review, we have summarised our understanding of the hydrology of High Mountain Asian
946 debris-covered glaciers, identified numerous knowledge gaps, and suggested six themes for future
947 research. While research has advanced substantially in recent years, there remain many questions
948 about how the hydrological systems of debris-covered glaciers behave, and how this varies
949 through both space and time. This limitation is largely due to the position of debris-covered
950 glaciers in hard-to-reach areas because of logistical difficulties and/or political instability, an
951 inability to gather observations beneath the surface due to the reduced performance of
952 combustion-powered equipment at high elevation, and the persistence of challenging weather
953 conditions for fieldwork through much of the year. Consequently, large uncertainty accompanies
954 any projections of future water supply, a concern for tens of millions of people across several
955 countries. Closing these knowledge gaps should thus prioritise generating information that best
956 improves robust model-based projections of water supply from High Mountain Asian debris-
957 covered glaciers. There is an inevitable trade-off between the cost of collecting the necessary
958 empirical data to close these gaps and the benefits returned in terms of improved model outputs.
959 In light of these requirements and considerations, we conclude by identifying two principal
960 priorities for scientists and two principal priorities for policymakers and funders.

961 Our first priority for scientists is to improve understanding of patterns and rates of surface
962 melting, particularly beneath debris layers of different properties and thicknesses on High
963 Mountain Asian debris-covered glaciers. To this end, multi-variable analytical models should be
964 developed to generate Østrem-type relationships applicable to a variety of debris properties (such
965 as lithology, shape, grain-size texture, and variability therein) and energy-balance regimes (thereby
966 factoring in influences such as elevation), extending the work of, for example Evatt et al. (2015).
967 Our second priority for scientists is to improve understanding of the basal hydrology of debris-
968 covered glaciers across all of High Mountain Asia. Currently, little is known about the subglacial
969 environment, including in many instances where the glacier base is, what the basal temperature
970 field is, and whether subglacial drainage occurs at all. Yet, these properties are central to the
971 quality of water supplied by such glaciers, as well as to their actual and modelled deformation
972 rates and motion fields, which govern their modelled response to anticipated climate change. In
973 order to maximise benefits relative to cost, field investigations of the subglacial environment could
974 be undertaken at a limited number of sites to evaluate and guide larger-scale remote sensing and
975 modelling studies.

976 Our first priority for policymakers and funders is to improve access for scientists to glaciers
977 across High Mountain Asia. In this regard, we believe the provision of a small number of bases with
978 effective transport infrastructure, open to international scientific teams, would facilitate a step
979 change in research activity and output. Our second priority for policymakers and funders is to
980 produce a better administrative environment for effective scientific collaboration. This should
981 include, for example, the development of memorandums of understanding between countries to
982 simplify regulations for research permitting and border crossing as part of a scientific research
983 project. It should also involve adopting best practice in terms of ensuring a uniform approach to
984 the quality control and homogeneity of data series, and archiving and sharing freely accessible

data. This would be in the interest of all involved parties, since maintaining a clean and reliable water supply is a fundamental part of building sustainable development (United Nations, 2015), which in High Mountain Asia can only be realised by improving understanding of future changes in the timing and magnitude of meltwater production from hitherto poorly studied debris-covered glaciers.

8. Author contributions

KM and BH planned the manuscript. KM led the manuscript writing and illustration. All authors contributed to the writing and editing of the manuscript.

9. Competing interests

The authors declare that they have no conflict of interest.

10. Acknowledgements

This research was supported by the ‘EverDrill’ Natural Environment Research Council Grant awarded to Aberystwyth University (NE/P002021) and the Universities of Leeds and Sheffield (NE/P00265X). KM is funded by an AberDoc PhD Studentship, with fieldwork costs supported by the Mount Everest Foundation and Postgraduate Research Awards from the British Society for Geomorphology, the Royal Geographical Society (with IBG) and Aberystwyth University Department of Geography and Earth Sciences. TIF acknowledges the Leverhulme Trust (RF-2018-584/4). The authors would like to thank Antony Smith (Aberystwyth University) for the initial illustration of Figure 2 and redrawing Figure 10. Figure 4 is reproduced from the Journal of Glaciology with permission of the International Glaciological Society. Figure 8 is reproduced according to a Creative Commons Attribution 4.0 License and Figure 10 is reproduced with permission of the publisher. They are also grateful to Himalayan Research Expeditions for organising the logistics that supported fieldwork in Nepal, and in particular Mahesh Magar for guiding and navigation. The authors thank two anonymous reviewers, and two previous reviewers in *The Cryosphere Discussions*, all the comments of which led to a much-improved manuscript.

11. References

- Adhikary, S., Nakawo, M., Seko, K., Shakya, B., 2000. Dust influence on the melting process of glacier ice: experimental results from Lirung Glacier, Nepal Himalayas, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 43–52.

- 1019 Ageta, Y., Iwata, S., Yabuki, H., Naito, N., Sakai, A., Narama, C., Karma, 2000. Expansion of glacier
1020 lakes in recent decades in the Bhutan Himalayas, in: Nakawo, M., Raymond, C.F., Fountain, A.
1021 (Eds.), *Debris-Covered Glaciers*. International Association of Hydrological Sciences, Oxford,
1022 pp. 165–175.
- 1023 Andermann, C., Bonnet, S., Crave, A., Davy, P., Longuevergne, L., Gloaguen, R., 2012a. Sediment
1024 transfer and the hydrological cycle of Himalayan rivers in Nepal. *Comptes Rendus Geosci.* 344,
1025 627–635. <https://doi.org/10.1016/j.crte.2012.10.009>
- 1026 Andermann, C., Crave, A., Gloaguen, R., Davy, P., Bonnet, S., 2012b. Connecting source and
1027 transport: Suspended sediments in the Nepal Himalayas. *Earth Planet. Sci. Lett.* 351–352,
1028 158–170. <https://doi.org/10.1016/j.epsl.2012.06.059>
- 1029 Andermann, C., Longuevergne, L., Bonnet, S., Crave, A., Davy, P., Gloaguen, R., 2012c. Impact of
1030 transient groundwater storage on the discharge of Himalayan rivers. *Nat. Geosci.* 5, 127–132.
1031 <https://doi.org/10.1038/ngeo1356>
- 1032 Anderson, R.S., 2000. A model of ablation-dominated medial moraines and the generation of
1033 debris-mantled glacier snouts. *J. Glaciol.* 46, 459–469.
- 1034 Barnett, T.P., Adam, J.C., Lettenmaier, D.P., 2005. Potential impacts of a warming climate on water
1035 availability in snow-dominated regions. *Nature* 438, 303–309.
1036 <https://doi.org/10.1038/nature04141>
- 1037 Basnett, S., Kulkarni, A.V., Bolch, T., 2013. The influence of debris cover and glacial lakes on the
1038 recession of glaciers in Sikkim Himalaya, India. *J. Glaciol.* 59, 1035–1046.
1039 <https://doi.org/10.3189/2013JoG12J184>
- 1040 Benn, D.I., Bolch, T., Hands, K., Gulley, J.D., Luckman, A., Nicholson, L., Quincey, D.J., Thompson,
1041 S.S., Toumi, R., Wiseman, S., 2012. Response of debris-covered glaciers in the Mount Everest
1042 region to recent warming, and implications for outburst flood hazards. *Earth-Science Rev.*
1043 114, 156–174. <https://doi.org/10.1016/j.earscirev.2012.03.008>
- 1044 Benn, D.I., Gulley, J.D., Luckman, A., Adamek, A., Glowacki, P.S., 2009. Englacial drainage systems
1045 formed by hydrologically driven crevasse propagation. *J. Glaciol.* 55, 513–523.
1046 <https://doi.org/10.3189/002214309788816669>
- 1047 Benn, D.I., Thompson, S.S., Gulley, J.D., Mertes, J.R., Luckman, A., Nicholson, L., 2017. Structure
1048 and evolution of the drainage system of a Himalayan debris-covered glacier, and its
1049 relationship with patterns of mass loss. *Cryosph.* 11, 2247–2264. [https://doi.org/10.5194/tc-](https://doi.org/10.5194/tc-2017-29)
1050 [2017-29](https://doi.org/10.5194/tc-2017-29)
- 1051 Benn, D.I., Wiseman, S., Hands, K.A., 2001. Growth and drainage of supraglacial lakes on debris-
1052 mantled Ngozumpa Glacier, Khumbu Himal, Nepal. *J. Glaciol.* 47, 626–638.
1053 <https://doi.org/10.3189/172756501781831729>
- 1054 Benn, D.I., Wiseman, S., Warren, C.R., 2000. Rapid growth of a supraglacial lake, Ngozumpa Glacier,
1055 Khumbu Himal, Nepal. *IAHS Publ.* 264, 177–185.
- 1056 Bhatt, M.P., Masuzawa, T., Yamamoto, M., Takeuchi, N., 2007. Chemical characteristics of pond
1057 waters within the debris area of Lirung Glacier in Nepal Himalaya. *J. Limnol.* 66, 71–80.
- 1058 Bocchiola, D., Senese, A., Mihalcea, C., Mosconi, B., D’Agata, C., Smiraglia, C., Diolaiuti, G., 2015.
1059 An ablation model for debris-covered ice: The case study of Venerocolo glacier (Italian Alps).
1060 *Geogr. Fis. e Din. Quat.* 38, 113–128. <https://doi.org/10.4461/GFDQ.2015.38.11>

1061 Bolch, T., 2017. Hydrology: Asian glaciers are a reliable water source. *Nature* 545, 161–162.
1062 <https://doi.org/10.1038/545161a>

1063 Bolch, T., Kulkarni, A., Kaab, A., Huggel, C., Paul, F., Cogley, J.G., Frey, H., Kargel, J.S., Fujita, K.,
1064 Scheel, M., Bajracharya, S., Stoffel, M., 2012. The State and Fate of Himalayan Glaciers.
1065 *Science*. 336, 310–314. <https://doi.org/10.1126/science.1215828>

1066 Bolch, T., Pieczonka, T., Benn, D.I., 2011. Multi-decadal mass loss of glaciers in the Everest area
1067 (Nepal Himalaya) derived from stereo imagery. *Cryosph.* 5, 349–358.
1068 <https://doi.org/10.5194/tc-5-349-2011>

1069 Bolch, T., Shea, J.M., Liu, S., Azam, F.M., Gao, Y., Gruber, S., 2019. Status and Change of the
1070 Cryosphere in the Extended Hindu Kush Himalaya Region, in: Wester, P., Mishra, A., Mukherji,
1071 A., Shrestha, A.B. (Eds.), *The Hindu Kush Himalaya Assessment*. Springer, Cham, Switzerland,
1072 pp. 209–255. https://doi.org/doi.org/10.1007/978-3-319-92288-1_7

1073 Bookhagen, B., Burbank, D.W., 2006. Topography, relief, and TRMM-derived rainfall variations
1074 along the Himalaya. *Geophys. Res. Lett.* 33, 1–5. <https://doi.org/10.1029/2006GL026037>

1075 Braun, L.N., Grabs, W., Rana, B., 1993. Application of a Conceptual Precipitation- Runoff Model in
1076 the Langtang Khola Basin , Nepal Himalaya. *Snow Glacier Hydrol. (Proceedings Kathmandu*
1077 *Symp. Novemb. 1992)*. IAHS Publ. no. 218,1993. 218, 221–237.

1078 Brock, B.W., Mihalcea, C., Kirkbride, M.P., Diolaiuti, G., Cutler, M.E.J., Smiraglia, C., 2010.
1079 Meteorology and surface energy fluxes in the 2005-2007 ablation seasons at the Miage
1080 debris-covered glacier, Mont Blanc Massif, Italian Alps. *J. Geophys. Res. Atmos.* 115, 1–16.
1081 <https://doi.org/10.1029/2009JD013224>

1082 Brun, F., Berthier, E., Wagnon, P., Käab, A., Treichler, D., 2017. A spatially resolved estimate of High
1083 Mountain Asia glacier mass balances from 2000 to 2016. *Nat. Geosci.* 10, 668–673.
1084 <https://doi.org/10.1038/ngeo2999>

1085 Brun, F., Buri, P., Miles, E.S., Wagnon, P., Steiner, J.F., Berthier, E., Ragettli, S., Kraaijenbrink, P.D.A.,
1086 Immerzeel, W.W., Pellicciotti, F., 2016. Quantifying volume loss from ice cliffs on debris-
1087 covered glaciers using high-resolution terrestrial and aerial photogrammetry. *J. Glaciol.* 62,
1088 684–695. <https://doi.org/10.1017/jog.2016.54>

1089 Brun, F., Wagnon, P., Berthier, E., Jomelli, V., Maharjan, S.B., Shrestha, F., Kraaijenbrink, P.D.A.,
1090 2019. Heterogeneous Influence of Glacier Morphology on the Mass Balance Variability in High
1091 Mountain Asia. *J. Geophys. Res. Earth Surf.* 124, 1331–1345.
1092 <https://doi.org/10.1029/2018JF004838>

1093 Brun, F., Wagnon, P., Berthier, E., Shea, J.M., Immerzeel, W.W., Kraaijenbrink, P.D.A., Vincent, C.,
1094 Reverchon, C., Shrestha, D., Arnaud, Y., 2018. Ice cliff contribution to the tongue-wide
1095 ablation of Changri Nup Glacier, Nepal, central Himalaya. *Cryosph.* 12, 3439–3457.
1096 <https://doi.org/10.5194/tc-12-3439-2018>

1097 Buri, P., Miles, E.S., Steiner, J.F., Immerzeel, W.W., Wagnon, P., Pellicciotti, F., 2016a. A physically-
1098 based 3-D model of ice cliff evolution on a debris-covered glacier. *J. Geophys. Res. Earth Surf.*
1099 121, 2471–2493. <https://doi.org/10.1002/2016JF004039>

1100 Buri, P., Pellicciotti, F., Steiner, J.F., Miles, E.S., Immerzeel, W.W., 2016b. A grid-based model of
1101 backwasting of supraglacial ice cliffs on debris-covered glaciers. *Ann. Glaciol.* 57, 199–211.
1102 <https://doi.org/10.3189/2016AoG71A059>

- 1103 Byers, A.C., Byers, E.A., McKinney, D.C., Rounce, D.R., 2017. A field-based study of impacts of the
1104 2015 earthquake on potentially dangerous glacial lakes in Nepal. *Himalaya* 37, 26–41.
- 1105 Carenzo, M., Pellicciotti, F., Mabillard, J., Reid, T., Brock, B.W., 2016. An enhanced temperature
1106 index model for debris-covered glaciers accounting for thickness effect. *Adv. Water Resour.*
1107 94, 457–469. <https://doi.org/10.1016/j.advwatres.2016.05.001>
- 1108 Carrivick, J.L., Tweed, F.S., 2013. Proglacial Lakes: Character, behaviour and geological importance.
1109 *Quat. Sci. Rev.* 78, 34–52. <https://doi.org/10.1016/j.quascirev.2013.07.028>
- 1110 Casey, K.A., Kääb, A., Benn, D.I., 2012. Geochemical characterization of supraglacial debris via in
1111 situ and optical remote sensing methods: a case study in Khumbu Himalaya, Nepal. *Cryosph.*
1112 6, 85–100. <https://doi.org/10.5194/tc-6-85-2012>
- 1113 Chaturvedi, R.K., Kulkarni, A., Karyakarte, Y., Joshi, J., Bala, G., 2014. Glacial mass balance changes
1114 in the Karakoram and Himalaya based on CMIP5 multi-model climate projections. *Clim.*
1115 *Change* 123, 315–328. <https://doi.org/10.1007/s10584-013-1052-5>
- 1116 Chen, C., Ding, Y., 2009. The application of artificial neural networks to simulate meltwater runoff
1117 of Keqikaer Glacier, south slope of Mt. Tuomuer, western China. *Environ. Geol.* 57, 1839–
1118 1845. <https://doi.org/10.1007/s00254-008-1471-1>
- 1119 Chikita, K., Jha, J., Yamada, T., 1998. The basin expansion mechanism of a supraglacial lake in the
1120 Nepal Himalaya. *J. Hokkaido Univ. Fac. Sci. Ser. VII Geophys.* 11, 501–521.
- 1121 Chu, V.W., 2014. Greenland ice sheet hydrology: A review. *Prog. Phys. Geogr.* 38, 19–54.
1122 <https://doi.org/10.1177/0309133313507075>
- 1123 Collier, E., Maussion, F., Nicholson, L., Mölg, T., Immerzeel, W.W., Bush, A., 2015. Impact of debris
1124 cover on glacier ablation and atmosphere-glacier feedbacks in the Karakoram. *Cryosph.* 9,
1125 1617–1632. <https://doi.org/10.5194/tc-9-1617-2015>
- 1126 Collier, E., Nicholson, L., Brock, B.W., Maussion, F., Essery, R.L.H., Bush, A.B.G., 2014. Representing
1127 moisture fluxes and phase changes in glacier debris cover using a reservoir approach.
1128 *Cryosph.* 8, 1429–1444. <https://doi.org/10.5194/tc-8-1429-2014>
- 1129 Conway, H., Rasmussen, L.A., 2000. Summer temperature profiles within supraglacial debris on
1130 Khumbu Glacier, Nepal, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered*
1131 *Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 89–97.
- 1132 Copland, L., Pope, S., Bishop, M.P., Shroder, J.F., Clendon, P., Bush, A., Kamp, U., Seong, Y.B., Owen,
1133 L.A., 2009. Glacier velocities across the Karakoram Himalaya. *Ann. Glaciol.* 50, 1–18.
1134 <https://doi.org/10.3189/172756409789624229>
- 1135 Dehecq, A., Gourmelen, N., Gardner, A.S., Brun, F., Goldberg, D., Nienow, P.W., Berthier, E.,
1136 Vincent, C., Wagnon, P., Trouvé, E., 2019. Twenty-first century glacier slowdown driven by
1137 mass loss in High Mountain Asia. *Nat. Geosci.* 12, 22–27. <https://doi.org/10.1038/s41561-018-0271-9>
- 1139 Dunning, S.A., Rosser, N.J., Mccoll, S.T., Reznichenko, N.V., 2015. Rapid sequestration of rock
1140 avalanche deposits within glaciers. *Nat. Commun.* 6, 1–7.
1141 <https://doi.org/10.1038/ncomms8964>
- 1142 Eriksson, M., Jianchu, X., Shrestha, A., Vaidya, R., Nepal, S., Sandström, K., 2009. The Changing
1143 Himalayas: Impact of climate change on water resources and livelihoods in the greater
1144 Himalayas. *Int. Cent. Integr. Mt. Dev.* 114, 1–28. <https://doi.org/10.1144/SP312.3>

- 1145 Evatt, G.W., Abrahams, I.D., Heil, M., Mayer, C., Kingslake, J., Mitchell, S.L., Fowler, A.C., Clark, C.D.,
1146 2015. Glacial melt under a porous debris layer. *J. Glaciol.* 61, 825–836.
1147 <https://doi.org/10.3189/2015JoG14J235>
- 1148 Fountain, A.G., Walder, J.S., 1998. Water flow through temperate glaciers. *Rev. Geophys.* 36, 299.
1149 <https://doi.org/10.1029/97RG03579>
- 1150 Fujita, K., Sakai, A., 2014. Modelling runoff from a Himalayan debris-covered glacier. *Hydrol. Earth*
1151 *Syst. Sci.* 18, 2679–2694. <https://doi.org/10.5194/hess-18-2679-2014>
- 1152 Fyffe, C.L., Brock, B.W., Kirkbride, M.P., Black, A.R., Smiraglia, C., Diolaiuti, G., 2019a. The influence
1153 of supraglacial debris on proglacial runoff fluctuations and water chemistry. *J. Hydrol.* 576,
1154 41–57. <https://doi.org/10.1016/j.jhydrol.2019.06.023>
- 1155 Fyffe, C.L., Brock, B.W., Kirkbride, M.P., Mair, D.W.F., Arnold, N.S., Smiraglia, C., Diolaiuti, G., Diotri,
1156 F., 2019b. Do debris-covered glaciers demonstrate distinctive hydrological behaviour
1157 compared to clean glaciers? *J. Hydrol.* 570, 584–597.
1158 <https://doi.org/10.1016/j.jhydrol.2018.12.069>
- 1159 Fyffe, C.L., Reid, T.D., Brock, B.W., Kirkbride, M.P., Diolaiuti, G., Smiraglia, C., Diotri, F., 2014. A
1160 distributed energy-balance melt model of an alpine debris-covered glacier. *J. Glaciol.* 60, 587–
1161 602. <https://doi.org/10.3189/2014JoG13J148>
- 1162 Gardelle, J., Arnaud, Y., Berthier, E., 2011. Contrasted evolution of glacial lakes along the Hindu
1163 Kush Himalaya mountain range between 1990 and 2009. *Glob. Planet. Change* 75, 47–55.
1164 <https://doi.org/10.1016/j.gloplacha.2010.10.003>
- 1165 Gardelle, J., Berthier, E., Arnaud, Y., 2012. Slight mass gain of Karakoram glaciers in the early
1166 twenty-first century. *Nat. Geosci.* 5, 322–325. <https://doi.org/10.1038/ngeo1450>
- 1167 Gibson, M.J., Glasser, N.F., Quincey, D.J., Mayer, C., Rowan, A.V., Irvine-Fynn, T.D.L., 2017a.
1168 Temporal variations in supraglacial debris distribution on Baltoro Glacier, Karakoram between
1169 2001 and 2012. *Geomorphology* 295, 572–585.
1170 <https://doi.org/10.1016/j.geomorph.2017.08.012>
- 1171 Gibson, M.J., Glasser, N.F., Quincey, D.J., Rowan, A.V., Irvine-Fynn, T.D.L., 2017b. Changes in glacier
1172 surface cover on Baltoro glacier, Karakoram, north Pakistan, 2001–2012. *J. Maps* 13, 100–108.
1173 <https://doi.org/10.1080/17445647.2016.1264319>
- 1174 Gilbert, A., Sinisalo, A., Gurung, T.R., Fujita, K., Maharjan, S.B., Sherpa, T.C., Fukuda, T., 2020. The
1175 influence of water percolation through crevasses on the thermal regime of a Himalayan
1176 mountain glacier. *Cryosph.* 14, 1273–1288. <https://doi.org/10.5194/tc-14-1273-2020>
- 1177 Gremaud, V., Goldscheider, N., Savoy, L., Favre, G., Masson, H., 2009. Geological structure,
1178 recharge processes and underground drainage of a glacierised karst aquifer system,
1179 Tsanfleuron-Sanetsch, Swiss Alps. *Hydrogeol. J.* 17, 1833–1848.
1180 <https://doi.org/10.1007/s10040-009-0485-4>
- 1181 Gulley, J.D., Benn, D.I., 2007. Structural control of englacial drainage systems in Himalayan debris-
1182 covered glaciers. *J. Glaciol.* 53, 399–412. <https://doi.org/10.3189/002214307783258378>
- 1183 Gulley, J.D., Benn, D.I., Müller, D., Luckman, A., 2009a. A cut-and-closure origin for englacial
1184 conduits in uncrevassed regions of polythermal glaciers. *J. Glaciol.* 55, 66–80.
1185 <https://doi.org/10.3189/002214309788608930>
- 1186 Gulley, J.D., Benn, D.I., Sreaton, E., Martin, J., 2009b. Mechanisms of englacial conduit formation

1187 and their implications for subglacial recharge. *Quat. Sci. Rev.* 28, 1984–1999.
 1188 <https://doi.org/10.1016/j.quascirev.2009.04.002>

1189 Hambrey, M.J., Quincey, D.J., Glasser, N.F., Reynolds, J.M., Richardson, S.J., Clemmens, S., 2008.
 1190 Sedimentological, geomorphological and dynamic context of debris-mantled glaciers, Mount
 1191 Everest (Sagarmatha) region, Nepal. *Quat. Sci. Rev.* 28, 1084.
 1192 <https://doi.org/10.1016/j.quascirev.2009.04.009>

1193 Han, H., Wang, J., Wei, J., Liu, S., 2010. Backwasting rate on debris-covered Koxkar glacier,
 1194 Tuomuer mountain, China. *J. Glaciol.* 56, 287–296.
 1195 <https://doi.org/10.3189/002214310791968430>

1196 Hannah, D.M., Kansakar, S.R., Gerrard, A.J., Rees, G., 2005. Flow regimes of Himalayan rivers of
 1197 Nepal: Nature and spatial patterns. *J. Hydrol.* 308, 18–32.
 1198 <https://doi.org/10.1016/j.jhydrol.2004.10.018>

1199 Haritashya, U.K., Kargel, J.S., Shugar, D.H., Leonard, G.J., Strattman, K., Watson, C.S., Shean, D.,
 1200 Harrison, S., Mandli, K.T., Regmi, D., 2018a. Evolution and controls of large glacial lakes in the
 1201 Nepal Himalaya. *Remote Sens.* 10, 1–31. <https://doi.org/10.3390/rs10050798>

1202 Haritashya, U.K., Kargel, J.S., Shugar, D.H., Leonard, G.J., Strattman, K., Watson, C.S., Shean, D.E.,
 1203 Harrison, S., Mandli, K.T., Regmi, D., 2018b. Evolution and controls of large glacial lakes in the
 1204 Nepal Himalaya. *Remote Sens.* 10, 1–31. <https://doi.org/10.3390/rs10050798>

1205 Haritashya, U.K., Kumar, A., Singh, P., 2010. Particle size characteristics of suspended sediment
 1206 transported in meltwater from the Gangotri Glacier, central Himalaya - An indicator of
 1207 subglacial sediment evacuation. *Geomorphology* 122, 140–152.
 1208 <https://doi.org/10.1016/j.geomorph.2010.06.006>

1209 Hasnain, S.I., 1999. Runoff characteristics of a glacierized catchment, Garhwal Himalaya, India.
 1210 *Hydrol. Sci. J.* 44, 847–854. <https://doi.org/10.1080/02626669909492284>

1211 Hasnain, S.I., 1996. Factors controlling suspended sediment transport in Himalayan glacier
 1212 meltwaters. *J. Hydrol.* 181, 49–62. [https://doi.org/10.1016/0022-1694\(95\)02917-6](https://doi.org/10.1016/0022-1694(95)02917-6)

1213 Hasnain, S.I., Jose, P.G., Ahmad, S., Negi, D.C., 2001. Character of the subglacial drainage system
 1214 in the ablation area of Dokriani glacier, India, as revealed by dye-tracer studies. *J. Hydrol.* 248,
 1215 216–223. [https://doi.org/10.1016/S0022-1694\(01\)00404-8](https://doi.org/10.1016/S0022-1694(01)00404-8)

1216 Hasnain, S.I., Thayyen, R.J., 1994. Hydrograph separation of bulk meltwaters of Dokriani Bamak
 1217 glacier basin, based on electrical conductivity. *Curr. Sci.* 67, 189–193.
 1218 <https://doi.org/24095811>

1219 Hewitt, K., Wake, C.P., Young, G.J., David, C., 1989. Hydrological investigations at Biafo Glacier,
 1220 Karakoram range, Himalaya: an important source of water for the Indus River. *Ann. Glaciol.*
 1221 13, 103–108. <https://doi.org/10.3189/S0260305500007710>

1222 Hock, R., Rasul, G., Adler, C., Caceres, S., Gruber, Y., Hirabayashi, Y., Jackson, M., Kääb, A., Kang, S.,
 1223 Kutuzov, S., Milner, A., Molau, U., Morin, S., Orlove, B., Steltzer, H., 2019. High Mountain
 1224 Areas, in: Pörtner, H.O., Roberts, D.C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska,
 1225 E., Mintenbeck, K., Nicolai, M., Okem, A., Petzold, J., Rama, B., Weyer, N. (Eds.), IPCC Special
 1226 Report on the Ocean and Cryosphere in a Changing Climate.

1227 Hodson, A.J., 2014. Understanding the dynamics of black carbon and associated contaminants in
 1228 glacial systems. *WIREs Water* 1, 141–149. <https://doi.org/10.1002/wat2.1016>

- 1229 Horodyskyj, U.N., 2015. Contributing factors to ice mass loss on Himalayan debris-covered glaciers.
1230 University of Colorado, Boulder.
- 1231 Hubbard, B., Nienow, P.W., 1997. Alpine subglacial hydrology. *Quat. Sci. Rev.* 16, 939–955.
1232 [https://doi.org/10.1016/S0277-3791\(97\)00031-0](https://doi.org/10.1016/S0277-3791(97)00031-0)
- 1233 Huss, M., 2011. Present and future contribution of glacier storage change to runoff from
1234 macroscale drainage basins in Europe. *Water Resour. Res.* 47, 1–14.
1235 <https://doi.org/10.1029/2010WR010299>
- 1236 Huss, M., Hock, R., 2018. Global-scale hydrological response to future glacier mass loss. *Nat. Clim.*
1237 *Chang.* 8, 135–140. <https://doi.org/10.1038/s41558-017-0049-x>
- 1238 Immerzeel, W.W., Kraaijenbrink, P.D.A., Shea, J.M., Shrestha, A.B., Pellicciotti, F., Bierkens, M.F.P.,
1239 De Jong, S.M., 2014. High-resolution monitoring of Himalayan glacier dynamics using
1240 unmanned aerial vehicles. *Remote Sens. Environ.* 150, 93–103.
1241 <https://doi.org/10.1016/j.rse.2014.04.025>
- 1242 Immerzeel, W.W., Lutz, A.F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S.,
1243 Davies, B.J., Elmore, A.C., Emmer, A., Feng, M., Fernandez, A., Haritashya, U., Kargel, J.S.,
1244 Hoppes, M., Kraaijenbrink, P.D.A., Kulkarni, A.V., Mayewski, P., Nepal, S., Pacheco, P., Painter,
1245 T.H., Pellicciotti, F., Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A.B., Viviroli, D., Wada, Y.,
1246 Xiao, C., Yao, T., Bailie, J.E.M., 2020. Importance and vulnerability of the world's water towers.
1247 *Nature* 577, 364–369. <https://doi.org/10.1038/s41586-019-1822-y>
- 1248 Immerzeel, W.W., van Beek, L.P.H., Bierkens, M.F.P., 2010. Climate change will affect the Asian
1249 water towers. *Science*. 328, 1382–1385. <https://doi.org/10.1126/science.1183188>
- 1250 Immerzeel, W.W., van Beek, L.P.H., Konz, M., Shrestha, A.B., Bierkens, M.F.P., 2012. Hydrological
1251 response to climate change in a glacierized catchment in the Himalayas. *Clim. Change* 110,
1252 721–736. <https://doi.org/10.1007/s10584-011-0143-4>
- 1253 Inoue, J., Yoshida, M., 1980. Ablation and Heat Exchange over the Khumbu Glacier. *J. Japanese Soc.*
1254 *Snow Ice* 41, 26–33. https://doi.org/10.5331/seppyo.41.Special_26
- 1255 Irvine-Fynn, T.D.L., Hodson, A.J., Moorman, B.J., Vatne, G., Hubbard, A.L., 2011. Polythermal
1256 Glacier Hydrology: A Review. *Rev. Geophys.* 49, 1–37.
1257 <https://doi.org/10.1029/2010RG000350>
- 1258 Irvine-Fynn, T.D.L., Porter, P.R., Rowan, A.V., Quincey, D.J., Gibson, M.J., Bridge, J.W., Watson, C.S.,
1259 Hubbard, A.L., Glasser, N.F., 2017. Supraglacial Ponds Regulate Runoff From Himalayan
1260 Debris-Covered Glaciers. *Geophys. Res. Lett.* 44, 11,894–11,904.
1261 <https://doi.org/10.1002/2017GL075398>
- 1262 Iwata, S., Watanabe, O., Fushimi, H., 1980. Surface morphology in the ablation area of the Khumbu
1263 Glacier. *J. Japanese Soc. Snow Ice* 41, 9–17. https://doi.org/10.5331/seppyo.41.Special_9
- 1264 Jansson, P., Hock, R., Schneider, T., 2003. The concept of glacier storage: a review. *J. Hydrol.* 282,
1265 116–129. [https://doi.org/10.1016/S0022-1694\(03\)00258-0](https://doi.org/10.1016/S0022-1694(03)00258-0)
- 1266 Jarosch, A.H., Gudmundsson, M.T., 2012. A numerical model for meltwater channel evolution in
1267 glaciers. *Cryosph.* 6, 493–503. <https://doi.org/10.5194/tc-6-493-2012>
- 1268 Juen, M., Mayer, C., Lambrecht, A., Han, H., Liu, S., 2014. Impact of varying debris cover thickness
1269 on ablation: A case study for Koxkar Glacier in the Tien Shan. *Cryosphere* 8, 377–386.
1270 <https://doi.org/10.5194/tc-8-377-2014>

- 1271 Kääb, A., 2005. Combination of SRTM3 and repeat ASTER data for deriving alpine glacier flow
1272 velocities in the Bhutan Himalaya. *Remote Sens. Environ.* 94, 463–474.
1273 <https://doi.org/10.1016/j.rse.2004.11.003>
- 1274 Kääb, A., Berthier, E., Nuth, C., Gardelle, J., Arnaud, Y., 2012. Contrasting patterns of early twenty-
1275 first-century glacier mass change in the Himalayas. *Nature* 488, 495–498.
1276 <https://doi.org/10.1038/nature11324>
- 1277 Kargel, J.S., Leonard, G.J., Shugar, D.H., Haritashya, U.K., Bevington, A., Fielding, E.J., Fujita, K.,
1278 Geertsema, M., Miles, E.S., Steiner, J., Anderson, E., Bajracharya, S., Bawden, G.W.,
1279 Breashears, D.F., Byers, A., Collins, B., Dhital, M.R., Donnellan, A., Evans, T.L., Geai, M.L.,
1280 Glasscoe, M.T., Green, D., Gurung, D.R., Heijenk, R., Hilborn, A., Hudnut, K., Huyck, C.,
1281 Immerzeel, W.W., Jiang, L., Jibson, R., Kääb, A., Khanal, N.R., Kirschbaum, D., Kraaijenbrink,
1282 P.D.A., Lamsal, D., Liu, S., Lv, M., McKinney, D., Nahirnick, N.K., Nan, Z., Ojha, S., Olsenholler,
1283 J., Painter, T.H., Pleasants, M., Pratima, K.C., Yuan, Q.I., Raup, B.H., Regmi, D., Rounce, D.R.,
1284 Sakai, A., Shangguan, D., Shea, J.M., Shrestha, A.B., Shukla, A., Stumm, D., Van Der Kooij, M.,
1285 Voss, K., Wang, X., Weihs, B., Wolfe, D., Wu, L., Yao, X., Yoder, M.R., Young, N., 2016.
1286 Geomorphic and geologic controls of geohazards induced by Nepal's 2015 Gorkha
1287 earthquake. *Science*. 351. <https://doi.org/10.1126/science.aac8353>
- 1288 Kattelmann, R., 2003. Glacial lake outburst floods in the Nepal Himalaya: A manageable hazard?
1289 *Nat. Hazards* 28, 145–154. <https://doi.org/10.1023/A:1021130101283>
- 1290 Kayastha, R.B., Takeuchi, Y., Nakawo, M., Ageta, Y., 2000. Practical prediction of ice melting
1291 beneath various thickness of debris cover on Khumbu Glacier, Nepal, using a positive degree-
1292 day factor, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered Glaciers*.
1293 International Association of Hydrological Sciences, Oxford, pp. 71–81.
- 1294 Kehrwald, N.M., Thompson, L.G., Tandong, Y., Mosley-Thompson, E., Schotterer, U., Alifimov, V.,
1295 Beer, J., Eikenberg, J., Davis, M.E., 2008. Mass loss on Himalayan glacier endangers water
1296 resources. *Geophys. Res. Lett.* 35, 2–7. <https://doi.org/10.1029/2008GL035556>
- 1297 Khan, A.A., Pant, N.C., Sarkar, A., Tandon, S.K., Thamban, M., Mahalinganathan, K., 2017. The
1298 Himalayan cryosphere: A critical assessment and evaluation of glacial melt fraction in the
1299 Bhagirathi basin. *Geosci. Front.* 8, 107–115. <https://doi.org/10.1016/j.gsf.2015.12.009>
- 1300 King, O., Bhattacharya, A., Bhambri, R., Bolch, T., 2019. Glacial lakes exacerbate Himalayan glacier
1301 mass loss. *Sci. Rep.* 9, 1–9. <https://doi.org/10.1038/s41598-019-53733-x>
- 1302 King, O., Quincey, D.J., Carrivick, J.L., Rowan, A.V., 2017. Spatial variability in mass loss of glaciers
1303 in the Everest region, central Himalaya, between 2000 and 2015. *Cryosph.* 11, 407–426.
1304 <https://doi.org/10.5194/tc-2016-99>
- 1305 Kirkbride, M.P., 2011. Debris-Covered Glaciers, in: Singh, V., Singh, P., Haritashya, U. (Eds.),
1306 *Encyclopedia of Snow, Ice and Glaciers*. Springer Netherlands, pp. 180–182.
1307 https://doi.org/10.1007/978-90-481-2642-2_622
- 1308 Kirkbride, M.P., 1993. The temporal significance of transitions from melting to calving termini in
1309 the glaciers of the central Southern Alps of New Zealand. *The Holocene* 3, 232–240.
- 1310 Kirkbride, M.P., Deline, P., 2013. The formation of supraglacial debris covers by primary dispersal
1311 from transverse englacial debris bands. *Earth Surf. Process. Landforms* 38, 1779–1792.
1312 <https://doi.org/10.1002/esp.3416>
- 1313 Kirkbride, M.P., Warren, C.R., 1999. Tasman Glacier, New Zealand: 20th-century thinning and

- 1314 predicted calving retreat. *Glob. Planet. Change* 22, 11–28. <https://doi.org/10.1016/S0921->
1315 8181(99)00021-1
- 1316 Kirkbride, M.P., Warren, C.R., 1997. Calving processes at a grounded ice cliff. *Ann. Glaciol.* 24, 116–
1317 121.
- 1318 Kodama, H., Mae, S., 1976. The Flow of Glaciers in the Khumbu Region. *J. Japanese Soc. Snow Ice*
1319 38, 31–36.
- 1320 Komori, J., 2008. Recent expansions of glacial lakes in the Bhutan Himalayas. *Quat. Int.* 184, 177–
1321 186. <https://doi.org/10.1016/j.quaint.2007.09.012>
- 1322 Kraaijenbrink, P.D.A., Bierkens, M.F.P., Lutz, A.F., Immerzeel, W.W., 2017. Impact of a global
1323 temperature rise of 1.5 degrees Celsius on Asia's glaciers. *Nature* 549, 257–260.
1324 <https://doi.org/10.1038/nature23878>
- 1325 Kraaijenbrink, P.D.A., Meijer, S.W., Shea, J.M., Pellicciotti, F., De Jong, S.M., Immerzeel, W.W.,
1326 2016a. Seasonal surface velocities of a Himalayan glacier derived by automated correlation
1327 of unmanned aerial vehicle imagery. *Ann. Glaciol.* 57, 103–113.
1328 <https://doi.org/10.3189/2016AoG71A072>
- 1329 Kraaijenbrink, P.D.A., Shea, J., Pellicciotti, F., de Jong, S., Immerzeel, W., 2016b. Object-based
1330 analysis of unmanned aerial vehicle imagery to map and characterise surface features on a
1331 debris-covered glacier. *Remote Sens. Environ.* 186, 581–595.
1332 <https://doi.org/10.1016/j.rse.2016.09.013>
- 1333 Kumar, K., Miral, M.S., Joshi, S., Pant, N., Joshi, V., Joshi, L.M., 2009. Solute dynamics of meltwater
1334 of Gangotri glacier, Garhwal Himalaya, India. *Environ. Geol.* 58, 1151–1159.
1335 <https://doi.org/10.1007/s00254-008-1592-6>
- 1336 Kumar, S., Dobhal, D.P., 1997. Climatic effects and bedrock control on rapid fluctuations of Chhota
1337 Shigri glacier, northwest Himalaya, India. *J. Glaciol.* 43, 467–472.
- 1338 Lejeune, Y., Bertrand, J., Wagnon, P., Morin, S., 2013. A physically based model of the year-round
1339 surface energy and mass balance of debris-covered glaciers. *J. Glaciol.* 59, 327–344.
1340 <https://doi.org/10.3189/2013JoG12J149>
- 1341 Li, J., Yuan, G., Wu, M., Sun, Y., Han, P., Wang, G., 2017. Evidence for persistent organic pollutants
1342 released from melting glacier in the central Tibetan Plateau, China. *Environ. Pollut.* 220, 178–
1343 185. <https://doi.org/10.1016/j.envpol.2016.09.037>
- 1344 Linsbauer, A., Frey, H., Haeberli, W., Machguth, H., Azam, M.F., Allen, S., 2016. Modelling glacier-
1345 bed overdeepenings and possible future lakes for the glaciers in the Himalaya-Karakoram
1346 region. *Ann. Glaciol.* 57, 119–130. <https://doi.org/10.3189/2016AoG71A627>
- 1347 Liu, Q., Liu, S., Cao, W., 2018. Seasonal Variation of Drainage System in the Lower Ablation Area of
1348 a Monsoonal Temperate Debris-Covered Glacier in Mt. Gongga, South-Eastern Tibet. *Water*
1349 10, 1–11. <https://doi.org/10.3390/w10081050>
- 1350 Liu, Q., Mayer, C., Liu, S., 2015. Distribution and interannual variability of supraglacial lakes on
1351 debris-covered glaciers in the Khan Tengri-Tumor Mountains, Central Asia. *Environ. Res. Lett.*
1352 10, 1–10. <https://doi.org/10.1088/1748-9326/10/1/014014>
- 1353 Lowe, A.T., Collins, D.N., 2001. Modelling runoff from large glacierized basins in the Karakoram
1354 Himalaya using remote sensing of the transient snowline. *Remote Sens. Hydrol.* 99–104.

1355 Lutz, A.F., Immerzeel, W.W., Shrestha, A.B., Bierkens, M.F.P., 2014. Consistent increase in High
1356 Asia's runoff due to increasing glacier melt and precipitation. *Nat. Clim. Chang.* 4, 587–592.
1357 <https://doi.org/10.1038/nclimate2237>

1358 Maharjan, S.B., Mool, P.K., Lizong, W., Xiao, G., Shrestha, F., Shrestha, R.B., Khanal, N.R.,
1359 Bajracharya, S.R., Joshi, S., Shai, S., Baral, P., 2018. The status of glacial lakes in the Hindu Kush
1360 Himalaya, ICIMOD Res. ed. ICIMOD, Kathmandu.

1361 Marston, R., 1983. Supraglacial Stream Dynamics on the Juneau Icefield. *Ann. Assoc. Am. Geogr.*
1362 73, 597–608.

1363 Mattson, L.E., 2000. The influence of a debris cover on the midsummer discharge of Dome Glacier,
1364 Canadian Rocky Mountains, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-
1365 Covered Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 25–33.

1366 Mayer, C., Lambrecht, A., Belò, M., Smiraglia, C., Diolaiuti, G., 2006. Glaciological characteristics of
1367 the ablation zone of Baltoro glacier, Karakoram, Pakistan. *Ann. Glaciol.* 43, 123–131.
1368 <https://doi.org/10.3189/172756406781812087>

1369 McCarthy, M., Pritchard, H., Willis, I.C., King, E., 2017. Ground-penetrating radar measurements of
1370 debris thickness on Lirung Glacier, Nepal. *J. Glaciol.* 63, 543–555.
1371 <https://doi.org/10.1017/jog.2017.18>

1372 Mertes, J.R., Thompson, S.S., Booth, A.D., Gulley, J.D., Benn, D.I., 2016. A conceptual model of
1373 supraglacial lake formation on debris-covered glaciers based on GPR facies analysis. *Earth
1374 Surf. Process. Landforms* 42, 903–914. <https://doi.org/10.1002/esp.4068>

1375 Mihalcea, C., Mayer, C., Diolaiuti, G., Lambrecht, A., Smiraglia, C., Tartari, G., 2006. Ice ablation
1376 and meteorological conditions on the debris-covered area of Baltoro glacier, Karakoram,
1377 Pakistan. *Ann. Glaciol.* 43, 292–300. <https://doi.org/10.3189/172756406781812104>

1378 Miles, E.S., Pellicciotti, F., Willis, I.C., Steiner, J.F., Buri, P., Arnold, N.S., 2016. Refined energy-
1379 balance modelling of a supraglacial pond, Langtang Khola, Nepal. *Ann. Glaciol.* 57, 29–40.
1380 <https://doi.org/10.3189/2016AoG71A421>

1381 Miles, E.S., Steiner, J.F., Willis, I.C., Buri, P., Immerzeel, W.W., Chesnokova, A., Pellicciotti, F.,
1382 2017a. Pond dynamics and supraglacial-englacial connectivity on debris-covered Lirung
1383 Glacier. *Front. Earth Sci.* 5. <https://doi.org/10.3389/feart.2017.00069>

1384 Miles, E.S., Watson, C.S., Brun, F., Berthier, E., Esteves, M., Quincey, D.J., Miles, K.E., Hubbard, B.,
1385 Wagnon, P., 2018a. Glacial and geomorphic effects of a supraglacial lake drainage and
1386 outburst event, Nepal Himalaya. *Cryosph.* 12, 3891–3905. [https://doi.org/10.5194/tc-2018-
1387 152](https://doi.org/10.5194/tc-2018-152)

1388 Miles, E.S., Willis, I.C., Arnold, N.S., Steiner, J.F., Pellicciotti, F., 2017b. Spatial, seasonal, and
1389 interannual variability of supraglacial ponds in the Langtang Valley of Nepal, 1999 to 2013. *J.
1390 Glaciol.* 63, 88–105. <https://doi.org/10.1017/jog.2016.120>

1391 Miles, E.S., Willis, I.C., Buri, P., Steiner, J.F., Arnold, N.S., Pellicciotti, F., 2018b. Surface pond energy
1392 absorption across four Himalayan glaciers accounts for 1/8 of total catchment ice loss.
1393 *Geophys. Res. Lett.* 45. <https://doi.org/10.1029/2018GL079678>

1394 Miles, K.E., Hubbard, B., Quincey, D.J., Miles, E.S., Irvine-Fynn, T.D.L., Rowan, A.V., 2019. Surface
1395 and subsurface hydrology of debris-covered Khumbu Glacier, Nepal, revealed by dye tracing.
1396 *Earth Planet. Sci. Lett.* 513, 176–186. <https://doi.org/doi.org/10.1016/j.epsl.2019.02.020>

- 1397 Miles, K.E., Hubbard, B., Quincey, D.J., Miles, E.S., Sherpa, T.C., Rowan, A.V., Doyle, S.H., 2018.
1398 Polythermal structure of a Himalayan debris-covered glacier revealed by borehole
1399 thermometry. *Sci. Rep.* 8, 1–9. <https://doi.org/10.1038/s41598-018-34327-5>
- 1400 Narama, C., Daiyrov, M., Tadono, T., Yamamoto, M., Kääh, A., Morita, R., Jinro, U., 2017. Seasonal
1401 drainage of supraglacial lakes on debris-covered glaciers in the Tien Shan Mountains, Central
1402 Asia. *Geomorphology* 286, 133–142. <https://doi.org/10.1016/j.geomorph.2017.03.002>
- 1403 Nicholson, L., Benn, D.I., 2013. Properties of natural supraglacial debris in relation to modelling
1404 sub-debris ice ablation. *Earth Surf. Process. Landforms* 38, 490–501.
1405 <https://doi.org/10.1002/esp.3299>
- 1406 Nicholson, L., Benn, D.I., 2006. Calculating ice melt beneath a debris layer using meteorological
1407 data. *J. Glaciol.* 52, 463–470. <https://doi.org/10.3189/172756506781828584>
- 1408 Nicholson, L.I., McCarthy, M., Pritchard, H.D., Willis, I., 2018. Supraglacial debris thickness
1409 variability: Impact on ablation and relation to terrain properties. *Cryosph.* 12, 3719–3734.
1410 <https://doi.org/10.5194/tc-12-3719-2018>
- 1411 Nie, Y., Sheng, Y., Liu, Q., Liu, L., Liu, S., Zhang, Y., Song, C., 2017. A regional-scale assessment of
1412 Himalayan glacial lake changes using satellite observations from 1990 to 2015. *Remote Sens.*
1413 *Environ.* 189, 1–13. <https://doi.org/10.1016/j.rse.2016.11.008>
- 1414 Nuimura, T., Fujita, K., Yamaguchi, S., Sharma, R.R., 2012. Elevation changes of glaciers revealed
1415 by multitemporal digital elevation models calibrated by GPS survey in the Khumbu region,
1416 Nepal Himalaya, 1992–2008. *J. Glaciol.* 58, 648–656. <https://doi.org/10.3189/2012JoG11J061>
- 1417 Østrem, G., 1959. Ice Melting under a Thin Layer of Moraine, and the Existence of Ice Cores in
1418 Moraine Ridges. *Geogr. Ann.* 41, 228–230.
1419 <https://doi.org/10.1080/20014422.1959.11907953>
- 1420 Pellicciotti, F., Stephan, C., Miles, E.S., Herreid, S., Immerzeel, W.W., Bolch, T., 2015. Mass-balance
1421 changes of the debris-covered glaciers in the Langtang Himal, Nepal, from 1974 to 1999. *J.*
1422 *Glaciol.* 61, 373–386. <https://doi.org/10.3189/2015JoG13J237>
- 1423 Pelto, M.S., 2000. Mass balance of adjacent debris-covered and clean glacier ice in the North
1424 Cascades, Washington, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered*
1425 *Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 35–42.
- 1426 Phillips, T., Rajaram, H., Steffen, K., 2010. Cryo-hydrologic warming: A potential mechanism for
1427 rapid thermal response of ice sheets. *Geophys. Res. Lett.* 37, 1–5.
1428 <https://doi.org/10.1029/2010GL044397>
- 1429 Pieczonka, T., Bolch, T., Junfeng, W., Shiyin, L., 2013. Heterogeneous mass loss of glaciers in the
1430 Aksu-Tarim Catchment (Central Tien Shan) revealed by 1976 KH-9 Hexagon and 2009 SPOT-5
1431 stereo imagery. *Remote Sens. Environ.* 130, 233–244.
1432 <https://doi.org/10.1016/j.rse.2012.11.020>
- 1433 Pohl, E., Gloaguen, R., Andermann, C., Knoche, M., 2017. Glacier melt buffers river runoff in the
1434 Pamir Mountains. *Water Resour. Res.* 53, 2467–2489. <https://doi.org/10.1002/2016WR019431>
- 1436 Pottakkal, J.G., Ramanathan, A., Singh, V.B., Sharma, P., Azam, M.F., Linda, A., 2014.
1437 Characterization of subglacial pathways draining two tributary meltwater streams through
1438 the lower ablation zone of Gangotri glacier system, Garhwal Himalaya, India. *Curr. Sci.* 107,

1439 613–621. <https://doi.org/24103533>

1440 Pritchard, H.D., 2019. Asia's shrinking glaciers protect large populations from drought stress.
1441 Nature 569, 649–654. <https://doi.org/10.1038/s41586-019-1240-1>

1442 Quincey, D.J., Braun, M., Glasser, N.F., Bishop, M.P., Hewitt, K., Luckman, A., 2011. Karakoram
1443 glacier surge dynamics. Geophys. Res. Lett. 38, 2–7. <https://doi.org/10.1029/2011GL049004>

1444 Quincey, D.J., Copland, L., Mayer, C., Bishop, M., Luckman, A., Belò, M., 2009. Ice velocity and
1445 climate variations for Baltoro Glacier, Pakistan. J. Glaciol. 55, 1061–1071.
1446 <https://doi.org/10.3189/002214309790794913>

1447 Quincey, D.J., Richardson, S.D., Luckman, A., Lucas, R.M., Reynolds, J.M., Hambrey, M.J., Glasser,
1448 N.F., 2007. Early recognition of glacial lake hazards in the Himalaya using remote sensing
1449 datasets. Glob. Planet. Change 56, 137–152.
1450 <https://doi.org/10.1016/j.gloplacha.2006.07.013>

1451 Radić, V., Bliss, A., Beedlow, A.C., Hock, R., Miles, E.S., Cogley, J.G., 2014. Regional and global
1452 projections of twenty-first century glacier mass changes in response to climate scenarios from
1453 global climate models. Clim. Dyn. 42, 37–58. <https://doi.org/10.1007/s00382-013-1719-7>

1454 Ragettli, S., Bolch, T., Pellicciotti, F., 2016a. Heterogeneous glacier thinning patterns over the last
1455 40 years in Langtang Himal, Nepal. Cryosph. 10, 2075–2097. <https://doi.org/10.5194/tc-10-2075-2016>

1457 Ragettli, S., Immerzeel, W.W., Pellicciotti, F., 2016b. Contrasting climate change impact on river
1458 flows from high-altitude catchments in the Himalayan and Andes Mountains. Proc. Natl. Acad.
1459 Sci. U. S. A. 113, 9222–9227. <https://doi.org/10.1073/pnas.1606526113>

1460 Ragettli, S., Pellicciotti, F., Immerzeel, W.W., Miles, E.S., Petersen, L., Heynen, M., Shea, J.M.,
1461 Stumm, D., Joshi, S., Shrestha, A., 2015. Unraveling the hydrology of a Himalayan catchment
1462 through integration of high resolution in situ data and remote sensing with an advanced
1463 simulation model. Adv. Water Resour. 78, 94–111.
1464 <https://doi.org/10.1016/j.advwatres.2015.01.013>

1465 Rana, B., Nakawo, M., Fukushima, Y., Ageta, Y., 1997. Application of a conceptual precipitation-
1466 runoff model (HYCYMODEL) in a debris-covered glacierized basin in the Langtang Valley,
1467 Nepal Himalaya. Ann. Glaciol. 25, 226–231.

1468 Raymond, C.F., Nolan, M., 2000. Drainage of a glacial lake through an ice spillway, in: Nakawo, M.,
1469 Raymond, C.F., Fountain, A. (Eds.), Symposium: Debris-Covered Glaciers. International
1470 Association of Hydrological Sciences, Oxford, pp. 199–210.

1471 Rees, G., Collins, D., 2006. Regional differences in response of flow in glacier-fed Himalayan rivers
1472 to climatic warming. Hydrol. Process. 20, 2157–2169. <https://doi.org/10.1002/hyp>

1473 Reid, T.D., Brock, B.W., 2014. Assessing ice-cliff backwasting and its contribution to total ablation
1474 of debris-covered Miage glacier, Mont Blanc massif, Italy. J. Glaciol. 60, 3–13.
1475 <https://doi.org/10.3189/2014JoG13J045>

1476 Reid, T.D., Brock, B.W., 2010. An energy-balance model for debris-covered glaciers including heat
1477 conduction through the debris layer. J. Glaciol. 56, 903–916.

1478 Reynolds, J.M., 2000. On the formation of supraglacial lakes on debris-covered glaciers. Debris-
1479 covered glaciers 264, 153–161.

- 1480 Röhl, K., 2008. Characteristics and evolution of supraglacial ponds on debris-covered Tasman
1481 Glacier, New Zealand. *J. Glaciol.* 54, 867–880. <https://doi.org/10.3189/002214308787779861>
- 1482 Röhl, K., 2006. Thermo-erosional notch development at fresh-water-calving Tasman Glacier, New
1483 Zealand. *J. Glaciol.* 52, 203–213. <https://doi.org/10.3189/172756506781828773>
- 1484 Rounce, D.R., Byers, A.C., Byers, E.A., McKinney, D.C., 2017. Brief Communications: Observations
1485 of a glacier outburst flood from Lhotse Glacier, Everest area, Nepal. *Cryosph.* 11, 443–449.
1486 <https://doi.org/10.5194/tc-2016-239>
- 1487 Rounce, D.R., Hock, R., Shean, D.E., 2020. Glacier Mass Change in High Mountain Asia Through
1488 2100 Using the Open-Source Python Glacier Evolution Model (PyGEM). *Front. Earth Sci.* 7, 1–
1489 20. <https://doi.org/10.3389/feart.2019.00331>
- 1490 Rounce, D.R., McKinney, D.C., Lala, J.M., Byers, A.C., Watson, C.S., 2016. A New Remote Hazard
1491 and Risk Assessment Framework for Glacial Lakes in the Nepal Himalaya. *Hydrol. Earth Syst.*
1492 *Sci.* 3455–3475. <https://doi.org/10.5194/hess-2016-161>
- 1493 Rowan, A.V., Egholm, D.L., Quincey, D.J., Glasser, N.F., 2015. Modelling the feedbacks between
1494 mass balance, ice flow and debris transport to predict the response to climate change of
1495 debris-covered glaciers in the Himalaya. *Earth Planet. Sci. Lett.* 430, 427–438.
1496 <https://doi.org/10.1016/j.epsl.2015.09.004>
- 1497 Sakai, A., 2012. Glacial Lakes in the Himalayas : A Review on Formation and Expansion Processes.
1498 *Glob. Environ. Res.* 16, 23–30.
- 1499 Sakai, A., Fujita, K., 2010. Formation conditions of supraglacial lakes on debris-covered glaciers in
1500 the Himalaya. *J. Glaciol.* 56, 9249–9251.
- 1501 Sakai, A., Nakawo, M., Fujita, K., 2002. Distribution Characteristics and Energy Balance of Ice Cliffs
1502 on Debris-Covered Glaciers, Nepal Himalaya. *Arctic, Antarct. Alp. Res.* 34, 12–19.
- 1503 Sakai, A., Nakawo, M., Fujita, K., 1998. Melt rate of ice cliffs on the Lirung Glacier, Nepal Himalayas,
1504 1996. *Bull. Glacier Res.* 16, 57–66.
- 1505 Sakai, A., Nishimura, K., Kadota, T., Takeuchi, N., 2009. Onset of calving at supraglacial lakes on
1506 debris-covered glaciers of the Nepal Himalaya. *J. Glaciol.* 55, 909–917.
1507 <https://doi.org/10.3189/002214309790152555>
- 1508 Sakai, A., Takeuchi, N., Fujita, K., Nakawo, M., 2000. Role of supraglacial ponds in the ablation
1509 process of a debris-covered glacier in the Nepal Himalayas, in: Nakawo, M., Raymond, C.F.,
1510 Fountain, A. (Eds.), *Debris-Covered Glaciers*. International Association of Hydrological
1511 Sciences, Oxford, pp. 119–130.
- 1512 Salerno, F., Guyennon, N., Thakuri, S., Viviano, G., Romano, E., Vuillermoz, E., Cristofanelli, P.,
1513 Stocchi, P., Agrillo, G., Ma, Y., Tartari, G., 2015. Weak precipitation, warm winters and springs
1514 impact glaciers of south slopes of Mt. Everest (central Himalaya) in the last 2 decades (1994–
1515 2013). *Cryosph.* 9, 1229–1247. <https://doi.org/10.5194/tc-9-1229-2015>
- 1516 Salerno, F., Thakuri, S., D’Agata, C., Smiraglia, C., Manfredi, E.C., Viviano, G., Tartari, G., 2012.
1517 Glacial lake distribution in the Mount Everest region: Uncertainty of measurement and
1518 conditions of formation. *Glob. Planet. Change* 92–93, 30–39.
1519 <https://doi.org/10.1016/j.gloplacha.2012.04.001>
- 1520 Savéan, M., Delclaux, F., Chevallier, P., Wagnon, P., Gonga-Saholiariliva, N., Sharma, R., Neppel, L.,
1521 Arnaud, Y., 2015. Water budget on the Dudh Koshi River (Nepal): Uncertainties on

1522 precipitation. *J. Hydrol.* 531, 850–862. <https://doi.org/10.1016/j.jhydrol.2015.10.040>

1523 Scherler, D., Bookhagen, B., Strecker, M.R., 2011. Spatially variable response of Himalayan glaciers
1524 to climate change affected by debris cover. *Nat. Geosci.* 4, 156–159.
1525 <https://doi.org/10.1038/ngeo1068>

1526 Scherler, D., Wulf, H., Gorelick, N., 2018. Global Assessment of Supraglacial Debris-Cover Extents.
1527 *Geophys. Res. Lett.* 45, 11,798–11,805. <https://doi.org/10.1029/2018GL080158>

1528 Scott, C.A., Zhang, F., Mukherji, A., Immerzeel, W.W., Mustafa, D., Bharati, L., 2019. Water in the
1529 Hindu Kush Himalaya, in: Wester, P., Mishra, A., Mukherji, A., Shrestha, A. (Eds.), *The Hindu*
1530 *Kush Himalaya Assessment*. Springer, Cham, Switzerland, pp. 257–299.
1531 https://doi.org/doi.org/10.1007/978-3-319-92288-1_8

1532 Shea, J.M., Immerzeel, W.W., 2016. An assessment of basin-scale glaciological and hydrological
1533 sensitivities in the Hindu Kush-Himalaya. *Ann. Glaciol.* 57, 308–318.
1534 <https://doi.org/10.3189/2016AoG71A073>

1535 Shea, J.M., Immerzeel, W.W., Wagnon, P., Vincent, C., Bajracharya, S., 2015. Modelling glacier
1536 change in the Everest region, Nepal Himalaya. *Cryosph.* 9, 1105–1128.
1537 <https://doi.org/10.5194/tc-9-1105-2015>

1538 Shean, D., 2017. High Mountain Asia 8-meter DEM Mosaics Derived from Optical Imagery, Version
1539 1. Tile 677. <https://doi.org/https://doi.org/10.5067/KXOVQ9L172S2>

1540 Shreve, R.L., 1972. Movement of Water in Glaciers. *J. Glaciol.* 11, 205–214.
1541 <https://doi.org/10.3189/S002214300002219X>

1542 Singh, P., Arora, M., Goel, N.K., 2006a. Effect of climate change on runoff of a glacierized Himalayan
1543 basin. *Hydrol. Process.* 20, 1979–1992. <https://doi.org/10.1002/hyp.5991>

1544 Singh, P., Bengtsson, L., 2004. Hydrological sensitivity of a large Himalayan basin to climate change.
1545 *Hydrol. Process.* 18, 2363–2385. <https://doi.org/10.1002/hyp.1468>

1546 Singh, P., Haritashya, U.K., Kumar, N., 2008. Modelling and estimation of different components of
1547 streamflow for Gangotri Glacier basin, Himalayas. *Hydrol. Sci. J.* 53, 309–322.
1548 <https://doi.org/10.1623/hysj.53.2.309>

1549 Singh, P., Haritashya, U.K., Kumar, N., Singh, Y., 2006b. Hydrological characteristics of the Gangotri
1550 Glacier, central Himalayas, India. *J. Hydrol.* 327, 55–67.
1551 <https://doi.org/10.1016/j.jhydrol.2005.11.060>

1552 Singh, P., Haritashya, U.K., Ramasastri, K.S., Kumar, N., 2005. Diurnal variations in discharge and
1553 suspended sediment concentration, including runoff-delaying characteristics, of the Gangotri
1554 Glacier in the Garhwal Himalayas. *Hydrol. Process.* 19, 1445–1457.
1555 <https://doi.org/10.1002/hyp.5583>

1556 Singh, P., Kumar, N., Ramasastri, K.S., Singh, Y., 2000. Influence of a fine debris layer on the melting
1557 of snow and ice on a Himalayan glacier, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.),
1558 *Debris-Covered Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 63–
1559 69.

1560 Singh, P., Ramasastri, K.S., Singh, U.K., Gergan, J.T., Dobhal, D.P., 1995. Hydrological characteristics
1561 of the Dokriani Glacier in the Garhwal Himalayas. *Hydrol. Sci. J.* 40, 243–257.
1562 <https://doi.org/10.1080/02626669509491407>

- Smart, C.C., 1996. Statistical evaluation of glacier boreholes as indicators of basal drainage systems. *Hydrol. Process.* 10, 599–613. [https://doi.org/10.1002/\(sici\)1099-1085\(199604\)10:4<599::aid-hyp394>3.0.co;2-8](https://doi.org/10.1002/(sici)1099-1085(199604)10:4<599::aid-hyp394>3.0.co;2-8)
- Smart, C.C., 1988. Artificial Tracer Techniques for the Determination of the Structure of Conduit Aquifers. *Groundwater*. <https://doi.org/10.1111/j.1745-6584.1988.tb00411.x>
- Smith, T., Bookhagen, B., 2018. Changes in seasonal snow water equivalent distribution in High Mountain Asia (1987 to 2009). *Sci. Adv.* 4. <https://doi.org/10.1126/sciadv.1701550>
- Soncini, A., Bocchiola, D., Confortola, G., Minora, U., Vuillermoz, E., Salerno, F., Viviano, G., Shrestha, D., Senese, A., Smiraglia, C., Diolaiuti, G., 2016. Future hydrological regimes and glacier cover in the Everest region: The case study of the upper Dudh Koshi basin. *Sci. Total Environ.* 1084–1101. <https://doi.org/10.1016/j.scitotenv.2016.05.138>
- Sorg, A., Bolch, T., Stoffel, M., Solomina, O., Beniston, M., 2012. Climate change impacts on glaciers and runoff in Tien Shan (Central Asia). *Nat. Clim. Chang.* 2, 725–731. <https://doi.org/10.1038/nclimate1592>
- Steiner, J.F., Buri, P., Miles, E.S., Ragettli, S., Pellicciotti, F., 2019. Supraglacial ice cliffs and ponds on debris-covered glaciers: Spatio-temporal distribution and characteristics. *J. Glaciol.* COMPLETE, 1–16. <https://doi.org/10.1017/jog.2019.40>
- Steiner, J.F., Kraaijenbrink, P.D.A., Jiduc, S.G., Immerzeel, W.W., 2018a. Brief communication: The Khurdopin glacier surge revisited - extreme flow velocities and formation of a dammed lake in 2017. *Cryosph.* 12, 95–101. <https://doi.org/10.5194/tc-12-95-2018>
- Steiner, J.F., Litt, M., Stigter, E.E., Shea, J.M., Bierkens, M.F.P., Immerzeel, W.W., 2018b. The Importance of Turbulent Fluxes in the Surface Energy Balance of a Debris-Covered Glacier in the Himalayas. *Front. Earth Sci.* 6, 1–18. <https://doi.org/10.3389/feart.2018.00144>
- Stigter, E.E., Litt, M., Steiner, J.F., Bonekamp, P.N.J., Shea, J.M., Bierkens, M.F.P., Immerzeel, W.W., 2018. The Importance of Snow Sublimation on a Himalayan Glacier. *Front. Earth Sci.* 6, 1–16. <https://doi.org/10.3389/feart.2018.00108>
- Stokes, C.R., Popovnin, V., Aleynikov, A., Gurney, S.D., Shahgedanova, M., 2007. Recent glacier retreat in the Caucasus Mountains, Russia, and associated increase in supraglacial debris cover and supra-/proglacial lake development. *Ann. Glaciol.* 46, 195–203. <https://doi.org/10.3189/172756407782871468>
- Sundal, A.V., Shepherd, A., Nienow, P.W., Hanna, E., Palmer, S., Huybrechts, P., 2009. Evolution of supra-glacial lakes across the Greenland Ice Sheet. *Remote Sens. Environ.* 113, 2164–2171. <https://doi.org/10.1016/j.rse.2009.05.018>
- Takeuchi, N., Sakai, A., Kohshima, S., Fujita, K., Nakawo, M., 2012. Variation in Suspended Sediment Concentration of Supraglacial Lakes on Debris-covered Area of the Lirung Glacier in the Nepal Himalayas. *Glob. Environ. Res.* 16, 95–104.
- Takeuchi, Y., Kayastha, R.B., Nakawo, M., 2000. Characteristics of ablation and heat balance in debris-free and debris-covered areas on Khumbu Glacier, Nepal Himalayas, in the pre-monsoon season, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 53–61.
- Tangborn, W., Rana, B., 2000. Mass balance and runoff of the partially debris- covered Langtang Glacier, Nepal, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered Glaciers*.

- 1605 International Association of Hydrological Sciences, Oxford, pp. 99–108.
- 1606 Thakuri, S., Salerno, F., Bolch, T., Guyennon, N., Tartari, G., 2016. Factors controlling the
1607 accelerated expansion of Imja Lake, Mount Everest region, Nepal. *Ann. Glaciol.* 57, 245–257.
1608 <https://doi.org/10.3189/2016AoG71A063>
- 1609 Thakuri, S., Salerno, F., Smiraglia, C., Bolch, T., D’Agata, C., Viviano, G., Tartari, G., 2014. Tracing
1610 glacier changes since the 1960s on the south slope of Mt. Everest (central Southern Himalaya)
1611 using optical satellite imagery. *Cryosph.* 8, 1297–1315. [https://doi.org/10.5194/tc-8-1297-](https://doi.org/10.5194/tc-8-1297-2014)
1612 2014
- 1613 Thayyen, R.J., Gergan, J.T., 2010. Role of glaciers in watershed hydrology : a preliminary study of a
1614 “Himalayan catchment.” *Cryosph.* 4, 115–128. <https://doi.org/10.5194/tcd-3-443-2009>
- 1615 Thayyen, R.J., Gergan, J.T., Dobhal, D.P., 2005. Monsoonal control on glacier discharge and
1616 hydrograph characteristics, a case study of Dokriani Glacier, Garhwal Himalaya, India. *J.*
1617 *Hydrol.* 306, 37–49. <https://doi.org/10.1016/j.jhydrol.2004.08.034>
- 1618 Thompson, S.S., Benn, D.I., Dennis, K., Luckman, A., 2012. A rapidly growing moraine-dammed
1619 glacial lake on Ngozumpa Glacier, Nepal. *Geomorphology* 145–146, 1–11.
1620 <https://doi.org/10.1016/j.geomorph.2011.08.015>
- 1621 Thompson, S.S., Benn, D.I., Mertes, J.R., Luckman, A., 2016. Stagnation and mass loss on a
1622 Himalayan debris-covered glacier: Processes, patterns and rates. *J. Glaciol.* 62, 467–485.
1623 <https://doi.org/10.1017/jog.2016.37>
- 1624 Tranter, M., Brown, G.H., Raiswell, R., Sharp, M.J., Gurnell, A., 1993. A conceptual model of solute
1625 acquisition by Alpine glacial meltwaters. *J. Glaciol.* 39, 573–581.
1626 <https://doi.org/10.3198/1993JoG39-133-573-581>
- 1627 Tranter, M., Sharp, M.J., Lamb, H.R., Brown, G.H., Hubbard, B., Willis, I.C., 2002. Geochemical
1628 weathering at the bed of Haut glacier d’Arolla, Switzerland - a new model. *Hydrol. Process.*
1629 16, 959–993. <https://doi.org/10.1002/hyp.309>
- 1630 Tsutaki, S., Fujita, K., Nuimura, T., Sakai, A., Sugiyama, S., Komori, J., 2019. Contrasting thinning
1631 patterns between lake- and land-terminating glaciers in the Bhutanese Himalaya. *Cryosph.*
1632 13, 2733–2750.
- 1633 United Nations, 2015. Transforming our world: the 2030 agenda for sustainable development,
1634 A/RES/70/1. <https://doi.org/10.1201/b20466-7>
- 1635 van Woerkom, T., Steiner, J.F., Kraaijenbrink, P.D.A., Miles, E.S., Immerzeel, W.W., 2019. Sediment
1636 supply from lateral moraines to a debris-covered glacier in the Himalaya. *Earth Surf. Dyn.* 7,
1637 411–427. <https://doi.org/10.5194/esurf-7-411-2019>
- 1638 Vaughan, D.G., Comiso, J.C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P., Murray, T., Paul,
1639 F., Ren, J., Rignot, E., Solomina, O., Steffen, K., Zhang, T., 2013. Observations: Cryosphere, in:
1640 Stocker, T.F., D. Qin, G.-K., Plattner, M., Tignor, S., Allen, K., Boschung, J., A.Nauels, Xia, Y.,
1641 Bex, V., Midgley, P.M. (Eds.), *Climate Change 2013: The Physical Science Basis. Contribution*
1642 *of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
1643 *Change.* Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp.
1644 317–382.
- 1645 Viste, E., Sorteberg, A., 2015. Snowfall in the Himalayas: an uncertain future from a little-known
1646 past. *Cryosph.* 9, 1147–1167. <https://doi.org/10.5194/tc-9-1147-2015>

- 1647 Wang, X., Ding, Y., Liu, S., Jiang, L., Wu, K., Jiang, Z., Guo, W., 2013. Changes of glacial lakes and
1648 implications in Tian Shan, central Asia, based on remote sensing data from 1990 to 2010.
1649 *Environ. Res. Lett.* 8, 044052. <https://doi.org/10.1088/1748-9326/8/4/044052>
- 1650 Wang, X., Liu, S., Han, H., Wang, J., Liu, Q., 2012. Thermal regime of a supraglacial lake on the
1651 debris-covered Koxkar Glacier, southwest Tianshan, China. *Environ. Earth Sci.* 67, 175–183.
1652 <https://doi.org/10.1007/s12665-011-1490-1>
- 1653 Watanabe, T., Lamsal, D., Ives, J.D., 2009. Evaluating the growth characteristics of a glacial lake
1654 and its degree of danger of outburst flooding: Imja Glacier, Khumbu Himal, Nepal. *Nor. Geogr.*
1655 *Tidsskr. - Nor. J. Geogr.* 63, 255–267. <https://doi.org/10.1080/00291950903368367>
- 1656 Watson, C.S., Quincey, D.J., Carrivick, J.L., Smith, M.W., 2017. Ice cliff dynamics in the Everest
1657 region of the Central Himalaya. *Geomorphology* 278, 238–251.
1658 <https://doi.org/10.1016/j.gloplacha.2016.04.008>
- 1659 Watson, C.S., Quincey, D.J., Carrivick, J.L., Smith, M.W., 2016. The dynamics of supraglacial ponds
1660 in the Everest region, central Himalaya. *Glob. Planet. Change* 142, 14–27.
1661 <https://doi.org/10.1016/j.gloplacha.2016.04.008>
- 1662 Watson, C.S., Quincey, D.J., Carrivick, J.L., Smith, M.W., Rowan, A.V., Richardson, R., 2018.
1663 Heterogeneous water storage and thermal regime of supraglacial ponds on debris-covered
1664 glaciers. *Earth Surf. Process. Landforms* 43, 229–241. <https://doi.org/10.1002/esp.4236>
- 1665 Wessels, R.L., Kargel, J.S., Kieffer, H.H., 2002. ASTER measurement of supraglacial lakes in the
1666 Mount Everest region of the Himalaya. *Ann. Glaciol.* 34, 399–408.
1667 <https://doi.org/10.3189/172756402781817545>
- 1668 Wilson, A.M., Williams, M.W., Kayastha, R.B., Racoviteanu, A., 2016. Use of a hydrologic mixing
1669 model to examine the roles of meltwater, precipitation and groundwater in the Langtang
1670 River basin, Nepal. *Ann. Glaciol.* 57, 155–168. <https://doi.org/10.3189/2016AoG71A067>
- 1671 Winiger, M., Gumpert, M., Yamout, H., 2005. Karakorum-Hindukush-western Himalaya: Assessing
1672 high-altitude water resources. *Hydrol. Process.* 19, 2329–2338.
1673 <https://doi.org/10.1002/hyp.5887>
- 1674 Zeng, C., Liu, Z., Yang, J., Yang, R., 2015. A groundwater conceptual model and karst-related carbon
1675 sink for a glacierized alpine karst aquifer, Southwestern China. *J. Hydrol.* 529, 120–133.
1676 <https://doi.org/10.1016/j.jhydrol.2015.07.027>
- 1677 Zhang, Y., Liu, S., Ding, Y., 2007. Glacier meltwater and runoff modelling, Keqicar Baqi glacier,
1678 southwestern Tien Shan, China. *J. Glaciol.* 53, 91–98.
1679 <https://doi.org/10.3189/172756507781833956>
- 1680 Zhao, L., Ding, R., Moore, J.C., 2014. Glacier volume and area change by 2050 in high mountain
1681 Asia. *Glob. Planet. Change* 122, 197–207. <https://doi.org/10.1016/j.gloplacha.2014.08.006>
- 1682